Canine tooth syndrome due to orbital roof fracture

Surabhi Dubey,¹ Oshin Bansal,² Ramesh Kekunnaya,³ Virender Sachdeva ⁴

¹Academy of Eye Care Education, Department of Pediatric Ophthalmology, Strabismus and Neuro-Ophthalmology, L V Prasad Eye Institute, GMR Varalakshmi Campus, Visakhapatnam, India

²Department of Ophthalmic Plastic Surgery, Ocular Oncology, Dacryology and Socket Sciences, L V Prasad Eye Institute, GMR Varalakshmi Campus, Visakhapatnam, India

³Child Sight Institute, Jasti V Ramanamma Children's Eye Care Centre, Department of Pediatric Ophthalmology, Strabismus and Neuro-ophthalmology, L V Prasad Eye Institute, KAR Campus, Hyderabad, Telangana, India

⁴Child Sight Institute, Nimaggada Prasad Children's Eye Care Centre, Paediatric Ophthalmology, Strabismus and Neuro-ophthalmology, L V Prasad Eye Institute, GMR Varalakshmi Campus, Visakhapatnam, Andhra Pradesh, India

Correspondence to

Dr Virender Sachdeva;
drvsachdeva29@gmail.com

Accepted 12 February 2022

DESCRIPTION

A young male presented to our clinic with complaints of binocular vertical double vision from 1 month. His complaints started following alleged motor vehicle accident with injury to the left side of the head. External examination showed a scar over the left forehead above the left eyebrow. His uncorrected visual acuity was 20/20, N6 in both eyes (BE), and colour vision was normal. Rest of the anterior segment examination was normal in BE.

Motor examination showed right head tilt of 10° (figure 1A). Hirschberg corneal reflex testing showed 5° of left hypertropia (LHT). Prism alternate cover testing for distance showed 6 prism diopters (PD) LHT with 5 PD exotropia (XT) in primary gaze. While XT was stable in all gazes, his LHT increased in the right gaze and on left head tilt to 9 PD. In the downgaze, his LHT increased to 14 PD. On extraocular motility testing, his ocular movements were full and free in the right eye (RE). However, the left eye (LE) showed –1 limitation of depression in adduction (dextrodepression). Additionally, he had –1 limitation of elevation which increased to –2 limitation of elevation in adduction (figure 1B). Fundus examination was normal in BE except for mild excyclotorsion in LE.

Based on the clinical findings, LHT worse in the right gaze and left head tilt, we diagnosed our patient to have superior oblique (SO) palsy (SOP) in the LE. However, our patient also had limitation of elevation, which was increasing in adduction, suggestive of associated Brown syndrome. We obtained CT scan of the orbit and brain which showed fracture of the medial wall extending to the medial part of the roof with injury to the trochlea and incarceration of SO at the site of the orbital fracture (figure 2A,B), leading to associated Brown syndrome-like presentation in our patient. We discussed the possible management options with oculoplastic colleagues, who advised exploration and reposition of the entrapped muscle. However, due

Figure 2 CT scan of the orbit on the coronal view shows the orbital roof fracture at the site of trochlea (A). On the axial view, CT scan shows entrapment of superior oblique muscle at the site of fracture (B).

to associated fracture of the anterior cranial fossa and a high risk of cerebrospinal fluid leak, the surgery was planned in coordination with neurosurgery department. However, the patient decided against surgery.

Our patient posed a diagnostic challenge and had features of both SOP and paradoxical limitation of elevation in adduction in the same eye which is classified as class 7 SOP according to Knapp's classification.¹ They described five cases of class 7 SOP (of which two had been associated with dog bite). Therefore, this entity is also termed as canine tooth syndrome. Other causes reported in the literature are due to surgical trauma to the trochlear region,² myocysticercosis of the SO³ and following closed head trauma.⁴ Adulkar *et al*⁵ reported a case of orbital roof fracture with SO entrapment at the site of fracture, where it resulted in acquired Brown syndrome-like presentation, but it did not have concomitant SO palsy. Therefore, our case would represent unique presentation of orbital roof fracture resulting in both SO palsy and acquired Brown syndrome.

Acknowledgements The authors would like to thank their Photography Department at L V Prasad Eye Institute, GMRV Campus, for all their support in the documentation and preparation of the images.

Contributors All authors contributed to the data collection, manuscript writing and final manuscript revision. VS conceptualised the idea.

Figure 1 Clinical photograph of the patient showing right head tilt and a depressed scar on the superior orbital rim at the site of the orbital fracture (yellow arrow). (A) In the primary gaze, the patient had left hypertropia of 7 prism diopters (PD) and 5 PD left exotropia. There is –1/2 limitation of depression which was worse in dextrodepression (black asterisk). (B) Further he had –1 limitation of elevation in primary gaze which increased to –2 limitation of elevation in adduction (small black arrow).

Learning points

- ▶ Fracture at the site of trochlea can cause both direct damage to the muscle and superior oblique (SO) palsy.
- ▶ Further entrapment of SO muscle at the fracture site can result in limitation of elevation in adduction, causing associated Brown syndrome and canine tooth-like syndrome.

© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Dubey S, Bansal O, Kekunnaya R, *et al.* *BMJ Case Rep* 2022;**15**:e248733. doi:10.1136/bcr-2021-248733

Funding This study was funded by Hyderabad Eye Research Foundation, Hyderabad, Telangana, India.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Virender Sachdeva <http://orcid.org/0000-0002-1927-6643>

REFERENCES

- 1 Knapp P, Moore S. Diagnosis and surgical options in superior oblique surgery. *Int Ophthalmol Clin* 1976;16:137–49.
- 2 Dikci S, Türkoğlu EB, Somer D. Canine tooth syndrome. *Turk J Ophthalmol* 2012;42:163–5.
- 3 Pandey PK, Bhatia A, Garg D, et al. Canine tooth syndrome due to superior oblique myocysticercosis. *J Pediatr Ophthalmol Strabismus* 2006;43:185–7.
- 4 Wong CW, Lim Z. A variant of canine tooth syndrome--presentation and management. *Strabismus* 2014;22:18–20.
- 5 Adulkar N, Kim U, Shetty S. Superior oblique muscle entrapment in orbital fracture presenting as acquired brown-like syndrome: a case report and review of literature. *Ophthalmic Plast Reconstr Surg* 2014;30:e26–8.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow