

Rare case of a large intracardiac serpiginous thrombus

Yixin Zhang ¹, Kevin Green,² Stuart Shah,³ Srinivasan Sattiraju⁴

¹Internal Medicine, University of Florida College of Medicine-Jacksonville, Jacksonville, Florida, USA

²Pulmonary and Critical Care Medicine, University of Florida Health at Jacksonville, Jacksonville, Florida, USA

³Pulmonary and Critical Care, University of Florida College of Medicine-Jacksonville, Jacksonville, Florida, USA

⁴Cardiology, University of Florida College of Medicine-Jacksonville, Jacksonville, Florida, USA

Correspondence to


Dr Yixin Zhang;
wendyz0411@gmail.com

Accepted 7 June 2022

DESCRIPTION

A man in his 50s presented to us with worsening fatigue and shortness of breath for 1 month. He recently underwent orthopaedics surgery and has been less active compared with baseline. Upon arrival, he was in obvious respiratory distress. Vital signs at the time showed blood pressure of 104/76 mm Hg, pulse rate of 90 beats per minute and pulse saturation of 92% on room air. The rest of the physical examination is unremarkable. The patient was put on 4 L of nasal cannula with symptomatic improvement. CT showed bilateral subsegmental pulmonary embolism (*figure 1*). Transthoracic echocardiogram (TTE) revealed a large serpiginous thrombus measuring at least 8 cm in the right atrium, intermittently protruding into the right ventricle (RV) along with severe RV dysfunction (*figure 2A, video 1*).

The patient was initially evaluated for a thrombectomy, but it was ultimately deferred due to the high motility of his intracardiac clot. We were worried that even gentle manipulation of the thrombus could possibly lead to secondary



Video 1 Off-axis apical four-chamber view showing a large serpiginous thrombus bouncing between the right atrium and right ventricle

embolism. Although he was haemodynamically stable at the time, we decided to administer 50 mg of intravenous systemic tissue plasminogen activator (tPA) infused over 2 hours due to the great embolic potential of his clot. Repeat





Figure 1 CT angiography of the chest showing bilateral subsegmental pulmonary embolism (red arrow).



Video 2 Off-axis apical four-chamber view taking 48 hours after tissue plasminogen activator administration showing complete resolution of the thrombus




Figure 2 (A) Off-axis apical four-chamber view showing a serpiginous clot between the right atrium (RA) and right ventricle (RV). (B) Complete resolution of the clot after administration of tissue plasminogen activator. LA, left atrium; LV, left ventricle.



© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Zhang Y, Green K, Shah S, et al. *BMJ Case Rep* 2022;**15**:e250545. doi:10.1136/bcr-2022-250545

Learning points

- ▶ Choosing the correct treatment modality in patients with intracardiac thrombus (eg, systemic thrombolytics, thrombectomy).
- ▶ Recognising possible fatal complications associated with a highly mobile intracardiac clot.
- ▶ The use of transthoracic echocardiography in thromboembolic disease can be lifesaving.

TTE 48 hours after tPA administration showed complete resolution of the clot (figure 2B, video 2). The patient also reported symptomatic relief without any further clinical deterioration. The patient was discharged with rivaroxaban after 5 days of hospital stay. This case provides striking TTE image and video of an exceptionally rare serpiginous clot moving freely between two cardiac chambers. It demonstrated that transthoracic echocardiography remains a valuable diagnostic modality for evaluation in thromboembolic disease and with timely use can be lifesaving.

Acknowledgements The authors acknowledge Hannah Burke, Sonographer for her contribution with echo images.

Contributors YZ drafted the manuscript and actively participated in patient care. KG proofread the final draft and actively participated in patient care. SS proofread

and edited the final draft and actively participated in patient care. SS drafted the manuscript and interpreted the diagnostic imaging.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Yixin Zhang <http://orcid.org/0000-0001-8154-0233>

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow