

Acute stroke presenting as syncope: Wallenberg syndrome

António Mesquita , Pedro Ferreira, Marta Carvalho, Miguel Martins

Serviço de Medicina 2.3,
Hospital Santo António dos
Capuchos, Lisboa, Portugal

Correspondence to
Dr António Mesquita;
aefpmesquita@gmail.com

Accepted 28 June 2022

DESCRIPTION

A man in his 50s presented to the emergency department with two episodes of syncope when standing up. They were not preceded by prodromal symptoms, chest pain, dyspnoea or palpitations. On arrival, the patient mentioned vertigo and instability when standing. Previous medical history was relevant for obesity, hypertension and liver transplant 3 years prior due to alcoholic cirrhosis. Physical examination showed multidirectional, non-suppressible, nystagmus, right-eye ptosis and myosis, loss of pain and temperature over the left side of the body (sparing face) and right-sided cerebellar hemiataxia. Romberg sign was absent due to severe unsteadiness when standing. Vital signs were normal and there was no difference between right-arm and left-arm pulse or blood pressure. Orthostatic vitals were not recorded. Blood glucose on admission was normal, as was a 12-lead ECG. CT of the head and neck with angiogram showed a hypoplastic left vertebral artery. Cranial MRI showed a T2 and fluid-attenuated inversion recovery (FLAIR) hyperintense lesion on the inferior lateral right bulbar region (figures 1 and 2), with restricted diffusion pattern (figure 3). Blood tests, including Rapid Plasma Reagin (RPR) and haemoglobin A1c levels, 24-hour Holter and transthoracic echocardiogram were normal. Doppler ultrasound of the cervical vessels was negative for subclavian steal syndrome or arterial dissection. Acute stroke from occlusion of the right posterior inferior cerebellar artery (PICA) was diagnosed. Clinical presentation was compatible with Wallenberg syndrome (WS). The patient showed almost complete resolution of his neurologic symptoms during admission.

Syncope is a form of transient and self-limited loss-of-consciousness caused by abrupt drop in cerebral blood flow, more specifically to the brainstem's reticular activating system. Major causes of syncope are reflex-mediated, such as vasovagal or carotid sinus hypersensitivity, orthostatic hypotension




Figure 2 Fluid-attenuated inversion recovery (FLAIR)-hyperintense lesion on the inferior lateral right medulla oblongata.

and cardiogenic¹. Although it results from brain hypoperfusion, stroke is only rarely the true cause of syncope, as the brain has a very redundant blood supply. One notable exception is the subclavian steal syndrome, which results in vertebrobasilar ischaemia during arm exercise due to subclavian artery stenosis. Stroke should be considered a cause of syncope when persistent and significant neurological changes occur before the episode or during the recovery period.

WS, also known as lateral medullary syndrome, is the most prevalent posterior ischaemic stroke syndrome. It commonly results from occlusion of the PICA or one of its branches. It is characterised by crossed thermoalgic hypoesthesia, ipsilateral cerebellar ataxia, vertigo, bulbar palsy and Horner's syndrome. Clinical presentation varies, often resulting in any combination of these symptoms. A wide range of unusual presentations of WS,




Figure 1 T2-hyperintense lesion on the inferior lateral right medulla oblongata, axial (A) and coronal (B) views.




Figure 3 Diffusion-weighted imaging with restricted diffusion (A) and low apparent diffusion coefficient (ADC) values (B) on the right lateral medulla oblongata.



© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Mesquita A, Ferreira P, Carvalho M, et al. *BMJ Case Rep* 2022;**15**:e251203. doi:10.1136/bcr-2022-251203

Table 1 Case reports of lateral medullary infarction with associated autonomic dysfunction

Case report	Manifestation of autonomic dysfunction
Huynh <i>et al</i> ⁴	Pre-syncope and orthostatic hypotension
Altavilla <i>et al</i> ⁵	Orthostatic hypotension and trigeminal-autonomic cephalalgia
Pascual <i>et al</i> ⁶	Retro-ocular headache with autonomic features resembling "continuous" cluster headache
Lee <i>et al</i> ⁷	Sick sinus syndrome
Takawaza <i>et al</i> ⁸	Sinus arrest and post-hiccup cough syncope
Alsaad <i>et al</i> ⁹	Sinus arrest and syncope, requiring permanent pacemaker placement

from trigeminal neuralgia to hiccups, has been reported in the literature.^{2,3} Autonomic dysfunction has been described in case reports, although rarely (table 1). Huynh *et al* reported a case of lateral medullary infarction presenting with presyncope in which autonomic testing showed impaired sympathetic and parasympathetic cardiovascular reflexes.⁴ Altavilla *et al* described a case of dorsolateral medullary infarction associated with orthostatic hypotension and trigeminal autonomic cephalalgia.⁵ Pascual and colleagues described another case presenting with cluster-like headache.⁶ In two separate case reports, sinus node dysfunction was diagnosed, with prolonged sinus pauses recorded on electrocardiography monitor.^{7,8} In another report, a patient presented with sinus arrest, requiring permanent pacemaker placement.⁹ To the best of our knowledge, however, only two cases of syncope associated with lateral medullary infarction have been reported, by Takawaza *et al* and Alsaad *et al*.^{8,9} These manifestations seem to be secondary to the involvement of sympathetic fibres and the *nucleus tractus solitarius* in the medulla, affecting the baroreceptor regulatory centre. In fact, Hong's study of autonomic

function in patients with lateral medullary infarction suggested susceptibility to cardiac parasympathetic dysfunction, especially in the ventral medulla.¹⁰

Our patient's syncope was compatible with orthostatic hypotension, which can be seen in autonomic failure. Although cardiac dysfunction following brainstem stroke is rare, this case highlights the importance of recognising syncope as a possible presenting symptom of medullary infarction and WS.

Contributors AM, PF, MC and MM contributed to manuscript elaboration and literature review.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s)

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

António Mesquita <http://orcid.org/0000-0003-3167-0990>

REFERENCES

- Shen W-K, Sheldon RS, Benditt DG, *et al*. 2017 ACC/AHA/HRS guideline for the evaluation and management of patients with syncope. *J Am Coll Cardiol* 2017;70:e39–110.
- Ordás CM, Cuadrado ML, Simal P, *et al*. Wallenberg's syndrome and symptomatic trigeminal neuralgia. *J Headache Pain* 2011;12:377–80.
- Sampath V, Gowda MR, Vinay HR, *et al*. Persistent hiccups (singultus) as the presenting symptom of lateral medullary syndrome. *Indian J Psychol Med* 2014;36:341–3.
- Huynh TR, Decker B, Fries TJ, *et al*. Lateral medullary infarction with cardiovascular autonomic dysfunction: an unusual presentation with review of the literature. *Clin Auton Res* 2018;28:569–76.
- Altavilla R, Landi D, Altamura C, *et al*. Dorsolateral medullary ischemic infarction causing autonomic dysfunction and headache: a case report. *J Headache Pain* 2012;13:327–30.
- Cid CG, Berciano J, Pascual J. Retro-ocular headache with autonomic features resembling "continuous" cluster headache in lateral medullary infarction. *J Neurol Neurosurg Psychiatry* 2000;69:134–5.
- Lee SW, Lee H, Yoo J, *et al*. Sick sinus syndrome combined with Wallenberg syndrome: a case report. *Brain Neurorehabil* 2021;14.
- Takawaza T, Ikeda K, Kano O, *et al*. A case of sinus arrest and post-hiccup cough syncope in medullary infarction. *J Stroke Cerebrovasc Dis* 2014;23:566–71.
- Alsaad AA, Austin CO, Robinson MT, *et al*. Pacemaker placement in patients with stroke-mediated autonomic dysregulation. *Case Rep Med* 2017;2017:1–3.
- Hong JM, Kim TJ, Shin DH, *et al*. Cardiovascular autonomic function in lateral medullary infarction. *Neurol Sci* 2013;34:1963–9.

Learning points

- ▶ Stroke is a rare cause of syncope.
- ▶ Persistent neurological deficits should prompt consideration of stroke as a cause of syncope.
- ▶ Syncope can be the presenting feature of Wallenberg syndrome, or lateral medullary infarction, secondary to autonomic dysfunction.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow