

# An atypical cause of headache

Mikel Vicente-Pascual <sup>1</sup>, Adriana Conde-Martín,<sup>2</sup> Angela Domenech-Cubí,<sup>2</sup> Cristina Molins-Rojas<sup>3</sup>

<sup>1</sup>Neurology Department, Xarxa Sanitària i Social de Santa Tecla, Tarragona, Catalunya, Spain

<sup>2</sup>General Medicine Department, Xarxa Sanitària i Social de Santa Tecla, Tarragona, Catalunya, Spain

<sup>3</sup>General Medicine Department, EAP Sant Pere i Sant Pau, Tarragona, Catalunya, Spain

## Correspondence to

Dr Mikel Vicente-Pascual;  
mvpscual.1@gmail.com

Accepted 2 September 2022

## DESCRIPTION

A woman in her 60s presented with wake-up right lateral cervical and occipital headache and gait instability. Six months ago, she suffered from a self-limiting episode of left-sided weakness, with a normal cranial CT. At the emergency department, she was haemodynamically stable. The clinical examination was normal, and the neurological exam showed gait instability, with no deficits in cranial nerves, muscular strength or sensibility. ECG, routine blood test and a new CT were normal. The ultrasound exam performed at the emergency department showed the right vertebral artery without flow (figure 1A), compatible with severe stenosis or occlusion, and with the normal flow in contralateral artery (figure 1B). Cerebral MRI with angiography showed an acute stroke in the right posterior inferior cerebellar artery territory and confirmed a filiform and discontinuous flow in the right vertebral artery (figure 2), compatible with artery dissection. She was treated with dual antiaggregation therapy without new symptoms.

Headaches are a common complaint in the emergency department. The main aetiology of headache is migraine, with arterial dissections being very rare.<sup>1</sup> The cervical artery dissection is a common cause of stroke in young adults. However, only 2% of strokes are due to dissections. One of the main symptoms is headache, which occurs in more than 70% of the patients,<sup>2</sup> followed by Horner's syndrome and cerebral or retinal ischaemia in carotid dissections, and ischaemia of the posterior circulation in vertebral dissection. The pain of vertebral artery dissection is commonly confused as musculoskeletal. To diagnose this entity, conventional arteriography is the gold standard. Our patient presented with a headache that wakes her up and neurological symptoms, both considered red flags. The



**Figure 2** MR angiography with filiform and discontinuous flow in the right vertebral artery, compatible with artery dissection.


localisation, the suddenness of the pain, the presence of gait instability and the previous paresis episode made us think of vascular aetiology, performing a sonography exam that was suggestive of severe stenosis or occlusion and an MRI with angiography that diagnosed the stroke and the vertebral dissection. In our opinion, this case shows the importance of vascular study in some types of headache.

## Learning points

- ▶ A vertebral dissection should be considered in patients with sudden unilateral headache or cervical pain.
- ▶ A CT cerebral angiogram should be performed if dissection is considered.

**Contributors** All authors contributed equally in data collection and data interpretation. AC-M, AD-C and MV-P contributed to patient management. MV-P designed the manuscript. MV-P and CM-R wrote the article. All authors read and approved the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.



**Figure 1** Supra-aortic trunk Doppler shows the right vertebral artery without flow (A), compatible with severe stenosis or occlusion, with normal flow in the contralateral artery (B).



© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Vicente-Pascual M, Conde-Martín A, Domenech-Cubí A, et al. *BMJ Case Rep* 2022;**15**:e252340. doi:10.1136/bcr-2022-252340

**Competing interests** None declared.

**Patient consent for publication** Consent obtained directly from patient(s).

**Provenance and peer review** Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

**ORCID iD**

Mikel Vicente-Pascual <http://orcid.org/0000-0002-5445-7853>

**REFERENCES**

- 1 Schievink WI. Spontaneous dissection of the carotid and vertebral arteries. *N Engl J Med Overseas Ed* 2001;344:898–906.
- 2 Vidale S. Headache in cervicocerebral artery dissection. *Neurol Sci* 2020;41:395–9.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

**Customer Service**

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at [support@bmj.com](mailto:support@bmj.com).

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow