



OPEN ACCESS

Experience of sudden sensorineural hearing loss following a mild COVID-19 infection

Kim Gibson,¹ John C C Ling,² Micah Peters³

¹Rosemary Bryant AO Research Centre, UniSA Clinical and Health Sciences, University of South Australia, Hawthorn, South Australia, Australia

²Specialists on Goodwood, Adelaide, South Australia, Australia

³Rosemary Bryant AO Research Centre, UniSA Clinical and Health Sciences, University of South Australia, Adelaide, South Australia, Australia

Correspondence to

Kim Gibson;
kim.gibson@unisa.edu.au

Accepted 15 March 2023

SUMMARY

A woman in her 30s was referred to an otolaryngologist with an acute onset of aural fullness, noise sensitivity, unilateral sudden onset hearing loss, vertigo and tinnitus. She had a confirmed COVID-19 infection 5 weeks prior. A pure tone audiogram confirmed sensorineural hearing loss. MRI identified an empty sella of the pituitary gland and without an obvious cause for hearing loss. Oral prednisolone and betahistine were prescribed, and her audiovestibular symptoms slowly improved over the subsequent months. The patient continues to experience intermittent tinnitus.

BACKGROUND

Sudden sensorineural hearing loss (SSNHL) is a syndrome characterised by rapid progression of hearing impairment (loss of 30 dB or more) over seconds to days.¹ In 84%–90% of cases, it is considered idiopathic and almost always unilateral.^{2,3} It is proposed that the SARS-CoV-2 virus attacks the auditory cortex in the temporal lobe, stimulating excess cytokine release. Cytokines then invade the cochlear nerve causing neuritis and the soft tissues of the cochlea causing cochleitis, leading to audiovestibular disturbance.⁴ Published evidence suggests potential association between recent upper respiratory tract infection and SSNHL,⁵ with increased prevalence reported during the COVID-19 pandemic.⁶ Some authors have suggested that SSNHL might be a symptom of COVID-19 and have observed the presence of SSNHL with or without other signs of infection.⁷

Most of the current literature on SSNHL and COVID-19 appear to be case studies that report development of SSNHL at different time points during an active COVID-19 infection with varying degrees of illness severity and audiological outcomes.^{8,9} There have also been reports of SSNHL arising following COVID-19 vaccination,¹⁰ however, evidence suggests that confirming an association between the two might be premature.¹¹ As evidence around the short-term and long-term impacts of COVID-19 and vaccines is still emerging and understanding rapidly evolving, it is important for clinicians to recognise both the variety of symptoms patients might present with as well as how these can be experienced by patients. With many reports of increasing burn-out among healthcare workers including cynicism/depersonalisation,^{12,13} acknowledging that even those who experience minor COVID-19 illness could be at risk of potentially permanent and debilitating long term effects is important to ensure high quality,

person-centred care.¹⁴ We report a case of an adult female (author and patient KG) who experienced SSNHL 5 weeks following a mild COVID-19 infection and continues to experience mild chronic tinnitus.

CASE PRESENTATION

After waking in the morning, a woman in her 30s first noted aural fullness, noise sensitivity and mild vertigo. Over the next 24 hours, these symptoms worsened, and she developed unilateral hearing loss and tinnitus in the right ear. She presented to her general practitioner and on otoscopic examination the ear canals appeared normal and non-inflamed with intact tympanic membranes. A referral to an otolaryngologist (one of the authors) was provided. Over the proceeding 24 hours, these symptoms worsened, and she further developed paroxysmal positional vertigo. She was reviewed by JCCL within 48 hours of her initial symptoms.

Relevant medical history included hypertension (managed with 4 mg perindopril daily) and resolved gestational diabetes (now 8 years post partum). She had received three doses of the BNT162b2 (Pfizer Comirnaty) COVID-19 vaccine. She had a polymerase chain reaction-confirmed COVID-19 infection 35 days prior to SSNHL onset. During her COVID-19 infection, she experienced sore throat, headache, nasal and sinus congestion, and lethargy. She did not require medical treatment and recovered by 2 weeks. She had no history of hearing loss or ear pathology and was relatively well prior to SSNHL but experienced ongoing mild nasal congestion since.

INVESTIGATIONS

Serial pure tone audiograms (PTA) are provided in figure 1. Forty eight hours after initial symptom development, a PTA was conducted measuring hearing thresholds from 250 Hz to 6000 Hz, identifying hearing loss in the right ear between 35 and 85 dB (figure 1A). MRI was conducted on the same day with no identifiable cause for acute right-sided hearing loss, but revealed an empty, slightly expanded sella of the pituitary gland, suggestive of a previous auto infarction of a possible pituitary adenoma.

PTA was repeated 3 weeks later indicating a reduction in hearing loss (30–70 dB loss between 250 and 6000 Hz) (figure 1B). An extended-frequency PTA (including air and bone conduction) was conducted the following month and hearing thresholds were measured from 250 Hz to 16 kHz (figure 1C). Hearing thresholds in the left ear were



© BMJ Publishing Group Limited 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gibson K, Ling JCC, Peters M. *BMJ Case Rep* 2023;**16**:e252899. doi:10.1136/bcr-2022-252899




Figure 1 Serial pure tone audiograms in sequential order, demonstrating gradual hearing improvement in the right ear. Blue, left ear; red, right ear. HL, hearing loss.

within normal limits for all frequencies tested, whereas a mild sensorineural hearing loss was observed in the right ear in the mid frequencies (500–3000 Hz) and extended high frequencies (8–16 kHz). A PTA at 3 months showed marked improvement with 10–35 dB loss (figure 1D). A final PTA at 4 months demonstrated hearing within normal limits (5–25 dB) (figure 1E).

TREATMENT

Audio-vestibular symptoms were treated with oral prednisone burst for 7 days (25 mg daily) and taper (12.5 mg daily) for another 7 days. A 12-day course of oral betahistine 16 mg twice daily was also prescribed.

OUTCOME AND FOLLOW-UP

Due to the empty sella of the pituitary gland on MRI, serum hormone levels were performed including thyroid stimulating hormone, follicle-stimulating hormone, luteinising hormone

and prolactin. All were reported to be within normal ranges and the patient is now under the care of an endocrinologist for ongoing surveillance. Three months since SSNHL, the patient now reports overall improvement but experiences intermittent pulsatile tinnitus.

DISCUSSION

This is an unusual case of SSNHL that developed 5 weeks following a mild COVID-19 infection. Like most cases of SSNHL where causation is difficult to prove, we postulate an association in this case especially since PTA revealed a sensory-neural mechanism for the unilateral hearing loss at the mid-to-high frequencies. Previous analysis of 20 asymptomatic people with COVID-19 showed deleterious effects on the cochlear hair cells and reduced amplitude of transient-evoked otoacoustic emissions compared with healthy individuals without a history of COVID-19.¹⁵

According to a recent systematic review and meta-analysis of cohort studies,¹⁶ during 2020–2021, approximately one-third of patients with SSNHL were positive for COVID-19 at the time of presentation to an audiology/otolaryngology referral centre. Moreover, in two included studies that reported epidemiological data, presentations of SSNHL increased during the pandemic.^{6 17} In a cross-sectional observational study by Dharmarajan *et al*,¹⁸ 100 patients who were hospitalised for COVID-19 received audiology testing that revealed sensorineural hearing loss in 53 of them.

There is heterogeneity in COVID-19 severity and the timing of audio-vestibular symptoms reported in published case studies. COVID-19 severity has ranged from asymptomatic or mild,^{7 9 19–26} moderate, requiring hospitalisation,²⁷ to severe, where intubation and intensive care were needed.^{28 29} Timing of audio-vestibular symptoms ranged from being the first indication of COVID-19^{7 19 30} to 3–12 days^{9 22 23} and between 4 and 12 weeks after reporting COVID-19 symptoms.^{20 21 25 27–29}

Based on the current evidence, the severity of COVID-19 infection and timing of SSNHL are therefore unpredictable. We recommend that clinicians apply a low threshold for suspicion of potential SSNHL among patients presenting with a wide variety of symptoms during the COVID-19 pandemic. While it is challenging to apprise patients regarding the sheer variety of potential short and long term COVID-19 impacts, we believe SSNHL should be included among the kinds of impacts clinicians consider discussing with patients. Rauch³ highlights that delayed diagnosis is a common challenge with cases of SSNHL with many

patients and clinicians attributing sensations of ‘ear fullness’ to impacted cerumen or congestion due to respiratory infection or allergies. Early administration of high-dose corticosteroids is a significant predictor of auditory outcome and is the recommend first-line treatment.³¹ It is therefore important for clinicians to carefully consider the potential for SSNHL and ensure prompt referral to specialists when symptoms first develop. This level of vigilance has also been highlighted by authors in relation to SSNHL occurring following COVID-19 infection.²⁵

SSNHL and associated symptoms can have a significant impact on quality of life,³² and are closely associated with anxiety and depression.^{33 34} This has implications for patient care and highlights the need for a holistic approach beyond medical treatment of SSNHL. Consideration of adjunct therapies such as stress reduction and well-being exercises may be worthwhile in tinnitus management and improvement in psychosocial outcomes.³⁵

In the coming months and years, it is likely that many currently unknown or poorly understood impacts of COVID-19 will be realised. As this area of research is largely limited to emerging studies with small population sizes, little is known about the risk factors for SSNHL to inform hearing screening protocols and early treatment with corticosteroids. It is vital that SSNHL be rigorously investigated among people who have active or recently recovered COVID-19 infections, as the risk of permanent audio-vestibular symptoms cannot be disregarded.

Twitter Kim Gibson @gibson_harvey

Acknowledgements The authors acknowledge the contribution of Boaz Mui, Flinders University.

Contributors KG: conceptualisation, writing-original draft preparation, patient. MP: conceptualisation, writing-original draft preparation, otolaryngologist. JCCL: draft review, expert opinion.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

Patient's perspective

Even working in health education, I had no previous knowledge of SSNHL being associated with COVID-19. This was a shocking experience for me that had a significant impact on my way of life for several months as I have never had any problems with hearing before; not even an ear infection. I was unable to drive a car while experiencing severe vertigo. I needed to reduce my workload, negotiate flexible working arrangements with my employer, and take a leave of absence from study. I felt uncomfortable and had difficulty conversing with people and attending social activities due to my hearing loss and noise sensitivity. I was worried that this would be permanent, and that I may require a hearing aid. I now feel nervous about a second COVID-19 infection. What if I experience this again or even worse, it results in permanent hearing loss?

Learning points

- ▶ Sudden sensorineural hearing loss (SSNHL) might be a symptom of COVID-19 infection and could also present weeks after full recovery even among patients who only experienced mild illness.
- ▶ The everyday impacts of SSNHL can be debilitating and distressing for patients and might lead to long term and potentially permanent disability.
- ▶ Clinicians should be aware of the variety of potential short-term and long-term impacts of COVID-19 and inform patients regarding what symptoms they should monitor for. Low thresholds for further investigation and follow-up should be considered for impacts that can potentially lead to long term or permanent disability.

REFERENCES

- 1 Chau JK, Lin JRJ, Atashband S, *et al*. Systematic review of the evidence for the etiology of adult sudden sensorineural hearing loss. *Laryngoscope* 2010;120:1011–21.
- 2 Kuhn M, Heman-Ackah SE, Shaikh JA, *et al*. Sudden sensorineural hearing loss: a review of diagnosis, treatment, and prognosis. *Trends Amplif* 2011;15:91–105.
- 3 Rauch SD. Clinical practice. idiopathic sudden sensorineural hearing loss. *N Engl J Med* 2008;359:833–40.
- 4 Harenberg J, Jonas JB, Trecca EMC. A liaison between sudden sensorineural hearing loss and SARS-cov-2 infection. *Thromb Haemost* 2020;120:1237–9.
- 5 Lin C-Y, Kuo P-H, Wu S-Y. Association of acute upper respiratory tract infections with sudden sensorineural hearing loss: a case-crossover, nationwide, population-based cohort study. *Int J Environ Res Public Health* 2021;18:20.
- 6 Fidan V, Akin O, Koyuncu H. Rised sudden sensorineural hearing loss during COVID-19 widespread. *Am J Otolaryngol* 2021;42:102996.
- 7 Kilic O, Kalcioğlu MT, Cag Y, *et al*. Could sudden sensorineural hearing loss be the sole manifestation of COVID-19? an investigation into SARS-COV-2 in the etiology of sudden sensorineural hearing loss. *Int J Infect Dis* 2020;97:208–11.
- 8 McIntyre KM, Favre NM, Kuo CC, *et al*. Systematic review of sensorineural hearing loss associated with COVID-19 infection. *Cureus* 2021;13:e19757.

- 9 Ricciardiello F, Pisani D, Viola P, *et al.* Sudden sensorineural hearing loss in mild COVID-19: case series and analysis of the literature. *Audiol Res* 2021;11:313–26.
- 10 Jeong J, Choi HS. Sudden sensorineural hearing loss after COVID-19 vaccination. *Int J Infect Dis* 2021;113:341–3.
- 11 Formeister EJ, Wu MJ, Chari DA, *et al.* Assessment of sudden sensorineural hearing loss after COVID-19 vaccination. *JAMA Otolaryngol Head Neck Surg* 2022;148:307–15.
- 12 Galanis P, Vraika I, Fragkou D, *et al.* Nurses' burnout and associated risk factors during the COVID-19 pandemic: a systematic review and meta-analysis. *J Adv Nurs* 2021;77:3286–302.
- 13 Morgantini LA, Naha U, Wang H, *et al.* Factors contributing to healthcare professional burnout during the COVID-19 pandemic: a rapid turnaround global survey. *PLoS ONE* 2020;15:e0238217.
- 14 Shanafelt TD. Enhancing meaning in work: a prescription for preventing physician burnout and promoting patient-centered care. *JAMA* 2009;302:1338–40.
- 15 Mustafa MWM. Audiological profile of asymptomatic covid-19 PCR-positive cases. *Am J Otolaryngol* 2020;41:102483.
- 16 Frosolini A, Franz L, Daloso A, *et al.* Sudden sensorineural hearing loss in the COVID-19 pandemic: a systematic review and meta-analysis. *Diagnostics (Basel)* 2022;12:12.
- 17 Parrino D, Frosolini A, Toninato D, *et al.* Sudden hearing loss and vestibular disorders during and before COVID-19 pandemic: an audiology tertiary referral centre experience. *Am J Otolaryngol* 2022;43:103241.
- 18 Dharmarajan S, Bharathi MB, Sivapuram K, *et al.* Hearing loss-a camouflaged manifestation of COVID 19 infection. *Indian J Otolaryngol Head Neck Surg* 2021;73:494–8.
- 19 Beckers E, Chouvel P, Cassetto V, *et al.* Sudden sensorineural hearing loss in COVID-19: a case report and literature review. *Clin Case Rep* 2021;9:2300–4.
- 20 Wong MY, Tang WS, Zakaria Z. Unilateral sudden sensorineural hearing loss in post-COVID-19 patients: case report. *Malays Fam Physician* 2022;17:112–6.
- 21 Pokharel S, Tamang S, Pokharel S, *et al.* Sudden sensorineural hearing loss in a post-COVID-19 patient. *Clin Case Rep* 2021;9:e04956.
- 22 Abdel Rhman S, Abdel Wahid A. COVID -19 and sudden sensorineural hearing loss, a case report. *Otolaryngol Case Rep* 2020;16:100198.
- 23 Edwards M, Muzaffar J, Naik P, *et al.* Catastrophic bilateral sudden sensorineural hearing loss following COVID-19. *BMJ Case Rep* 2021;14:e243157.
- 24 Ozer F, Alkan O. n.d. Simultaneous sudden hearing loss and peripheral facial paralysis in a patient with covid-19. *Ear Nose Throat J*;2021:014556132110280.
- 25 Lang B, Hintze J, Conlon B. Coronavirus disease 2019 and sudden sensorineural hearing loss. *J Laryngol Otol* 2020;134:1–3.
- 26 Karimi-Galougahi M, Naeini AS, Raad N, *et al.* Vertigo and hearing loss during the COVID-19 pandemic-is there an association? *Acta Otorhinolaryngol Ital* 2020;40:463–5.
- 27 Chakraborty S, Maharatna S, Reddy CS, *et al.* Hearing loss in a post COVID-19 patient. *Int J Otorhinolaryngol Head Neck Surg* 2021;7:548.
- 28 Koumpa FS, Forde CT, Manjaly JG. Sudden irreversible hearing loss post COVID-19. *BMJ Case Rep* 2020;13:11.
- 29 Gerstaecker K, Speck I, Riemann S, *et al.* Deafness after COVID-19? *HNO* 2021;69(Suppl 2):92–5.
- 30 Chern A, Famuyide AO, Moonis G, *et al.* Bilateral sudden sensorineural hearing loss and intralabyrinthine hemorrhage in a patient with COVID-19. *Otol Neurotol* 2021;42:e10–4.
- 31 Foden N, Mehta N, Joseph T. Sudden onset hearing loss -- causes, investigations and management. *Aust Fam Physician* 2013;42:641–4.
- 32 Saniasaya J. Hearing loss in SARS-cov-2: what do we know? *Ear Nose Throat J* 2021;100(2_suppl):152S–154S.
- 33 Trevis KJ, McLachlan NM, Wilson SJ. A systematic review and meta-analysis of psychological functioning in chronic tinnitus. *Clin Psychol Rev* 2018;60:62–86.
- 34 Bhatt JM, Bhattacharyya N, Lin HW. Relationships between tinnitus and the prevalence of anxiety and depression. *Laryngoscope* 2017;127:466–9.
- 35 Gans JJ, O'Sullivan P, Bircheff V. Mindfulness based tinnitus stress reduction pilot study: A symptom perception-shift program. *Mindfulness (N Y)* 2013;5:322–33.

Copyright 2023 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow