

Non-traumatic parasternal pericardiocentesis

Pitt O Lim 

Department of Cardiology,
Atkinson-Morley Wing, St
George's University Hospitals
NHS Foundation Trust, Tooting,
UK

Correspondence to

Dr Pitt O Lim;
pitt.lim@stgeorges.nhs.uk

Accepted 13 April 2023

DESCRIPTION

A breathless woman in her 30s was admitted with cardiac tamponade following resection of her left upper lung lobe with carcinoid tumour over 6 months prior. She had inflammatory polyarthropathy and incidental pulmonary embolism confirmed on CT pulmonary angiogram. This was treated with full anticoagulation with low-molecular-weight heparin. To perform her pericardiocentesis, three possible methods were considered; (1) subcostal approach had overlying liver and abdominal adiposity rendering this conventional access inadvisable, published series indicate an up to 20% complications rate which is also the most indirect route and in current practice it accounts for the bulk of adverse events^{1,2}; (2) apical approach was hindered by breast, chest adiposity and overlying lung with the potential of causing pneumothorax and finally; (3) parasternal approach which is increasingly done in the emergency department by non-cardiologists and was favourable.³ Her echocardiogram (figure 1A) revealed significant effusion around the heart with tamponade physiology on doppler and evidence of diastolic right ventricular

collapse. The ultrasound probe was used to mark an ideal "X" spot 1cm from the left sternal edge in the fourth intercostal space with the patient in supine posture, overlying the cardiac notch. This area is usually devoid of lung tissue, where the parietal pericardium abuts the chest wall and this measured 3cm for this patient (figure 1B). Importantly, the left internal mammary artery which runs > 13mm from the sternum could be avoided. A 21 gauge needle from the Cook Micropuncture Pedal Introducer Access Set (Cook Medical, Bloomington, USA) was used to enter the pericardial space; this needle has a length of 4 cm, which means that there is no risk of going too deep to penetrate any cardiac structures. Therefore removing the need for simultaneous ultrasound guidance during needling. Otherwise, using a bigger and longer 18 gauge x 15 cm standard needle, this would have to be advanced at an angulation under the ultrasound probe with the potential of puncturing the left internal mammary artery.^{3,4} In this case, as the needle was inserted vertically, the first flush-back indicated entry into the pericardial space and a thin 0.018" wire was placed as seen in the fluoroscopic




Figure 1 Large pericardial effusion on echocardiogram (A), distance between chest wall and parietal pericardium (B, arrow), small 21 gauge needle and thin 0.018" wire in pericardial space (C) and pericardial drain in situ (D).



© BMJ Publishing Group
Limited 2023. No commercial
re-use. See rights and
permissions. Published by BMJ.

To cite: Lim PO. *BMJ Case Rep* 2023;**16**:e253728.
doi:10.1136/bcr-2022-253728

Learning points

- Pericardial effusion causing cardiac tamponade is an increasingly common medical emergency. Recent general availability of echocardiogram in the accident and emergency department means that the sickest patients with it are being diagnosed requiring on-the-spot pericardiocentesis.
- There are three common approaches to pericardiocentesis: subcostal, apical and parasternal. The first approach though conventional and standard practice is the least direct and is associated with the most complications.
- Applying a slender technique to the parasternal approach to pericardiocentesis potentially makes the procedure simple, safe and predictable.

picture in left anterior oblique (LAO) view (figure 1C). Utilising the Seldinger Method, a 4 French cannula (diameter 1.33 mm × length 7 cm) which came in the pack was put into the pericardial space for pressure measurements. The mean pericardial pressure was 17 (< 4) mm Hg, confirming cardiac tamponade. Finally, a 0.035" wire from the PERIVAC Pericardiocentesis Kit (Boston Scientific, Massachusetts, USA) was introduced via the cannula and an 8.3 French drain provided was positioned with ease along the upper border of the rib thus avoiding the neurovascular bundle, as seen in the lower fluoroscopic picture in posterior anterior (PA) view (figure 1D). After 200 mL of blood-stained fluid was withdrawn, the pericardial pressure fell to a mean of 0 mmHg with immediate symptomatic relief. Slender techniques are generally non-traumatic.⁵ This two-step technique was previously described elsewhere and it allows accessing the pericardial space at a first pass in a fully anticoagulated patient.⁶ This additional step potentially renders the procedure as safe as a diagnostic pleural tap.

Acknowledgements The author thanks Drs Rajan Sharma and Manav Sohal (clinical leaders), Mary Keal and Menene Endaya (cath lab nursing), Dinesh Sajani and Jaysson Crusis (cardiac radiography) and Claire Oneill (cardiac physiology); Matthew Li Kam Wa (trainee) who assisted in the case; Departmental and Cardiac Catheter Laboratory auxiliary and nursing staff involved in caring for the patient; the patient who graciously consented to having her case shared for medical learning and communication; and Farida Ismail for her meticulous stock keeping.

Contributors POL assessed the patient, performed the procedure and followed up the patient in his outpatient clinic. He wrote the manuscript and approved it for publication.

Funding The author has not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Pitt O Lim <http://orcid.org/0000-0001-8821-3008>

REFERENCES

- 1 Stolz L, Situ-LaCasse E, Acuña J, *et al.* What is the ideal approach for emergent pericardiocentesis using point-of-care ultrasound guidance? *World J Emerg Med* 2021;12:169–73.
- 2 Inglis R, King AJ, Gleave M, *et al.* Pericardiocentesis in contemporary practice. *J Invasive Cardiol* 2011;23:234–9.
- 3 Osman A, Wan Chuan T, Ab Rahman J, *et al.* Ultrasound-Guided pericardiocentesis: a novel parasternal approach. *Eur J Emerg Med* 2018;25:322–7.
- 4 Ferro B, Vegnuti L, Santonocito O, *et al.* Ultrasound emergency lateral-to-medial parasternal pericardiocentesis with high frequency probe in COVID-19: a case report. *Eur Heart J Case Rep* 2022;6:ytac203.
- 5 Lim PO, Elghamry Z. Heparin-free distal radial artery approach to cardiac catheterisation and the small radial recurrent artery. *Br J Cardiol* 2021;28:39:39.
- 6 Ferreira-Martins J, Lim PO. Two-Step distal radial artery cannulation for challenging radial anatomies. *Cardiology* 2021;146:144–50.

Copyright 2023 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow