

Scrub typhus: the clinical significance of the eschar

Takashi Akaike,¹ Kosuke Ishizuka ,² Naoto Tominaga,³ Iori Motohashi¹

¹Division of General Internal Medicine, Department of Internal Medicine, Kawasaki Municipal Tama Hospital, Kawasaki, Kanagawa, Japan

²Department of General Medicine, Yokohama City University School of Medicine, Yokohama, Kanagawa, Japan

³Division of Nephrology and Hypertension, Department of Internal Medicine, Kawasaki Municipal Tama Hospital, Kawasaki, Kanagawa, Japan

Correspondence to

Dr Kosuke Ishizuka;
e103007c@yokohama-cu.ac.jp

Accepted 18 April 2023

DESCRIPTION

A man in his 80s on chronic haemodialysis for end-stage kidney disease due to microscopic polyangiitis (MPA) presented with fever and chills since the morning. His medications included prednisolone (5 mg/day) and azathioprine (50 mg/day) for MPA. He had also engaged in leisure gardening. Physical examination showed a body temperature of 39.1°C, heart rate of 100 beats per minute, blood pressure of 118/65 mm Hg, and respiratory rate of 20 breaths per minute. Laboratory data exhibited the following: aspartate aminotransferase, 86 U/L (reference range: 13–30 U/L); alanine aminotransferase, 35 U/L (reference range: 10–42 U/L); C reactive protein, 8.03 mg/dL (reference range: 0.00–0.14 mg/dL); WBC, 3800/μL (neutrophils, 96.0%; lymphocytes, 2.5%; reference range: 4000–9000/μL); platelets, 42 000/μL (reference range: 158 000–348 000/μL); and myeloperoxidase antineutrophil cytoplasmic antibody; 1.1 IU/L (reference range <3.5 IU/L). Detailed physical examination revealed an eschar of 5 mm diameter on the medial side of the right lower leg ([figure 1](#)). PCR of an eschar biopsy specimen was positive for *Orientia tsutsugamushi*, and scrub typhus was diagnosed. His symptoms resolved after treatment with minocycline, 200 mg/day for 7 days.

Scrub typhus, a zoonotic disease also known as tsutsugamushi disease, is a rickettsiosis caused by *Orientia tsutsugamushi*, which is frequently reported in Asia and the Pacific Rim.¹ Although it varies by region, tsutsugamushi disease peaks in the fall and early winter.² The disease is transmitted by the bite of a tick larva, called chigger, and after an incubation period of 7–10 days, symptoms such as headache, myalgia

and rash are observed besides fever and chills. While the accompanying symptoms other than fever and chills may be unremarkable, they are often fatal.³ As seen in this case, an infiltrative erythema with a 5–15 mm long black eschar at the site of the infected chigger bite can be often found on physical examination. Approximately 80% of patients have an eschar, which is of high diagnostic value, but patients themselves do not often notice the presence of the eschar, which may be found in obscure sites such as the pubis, axilla or lower legs; therefore, careful physical examination is necessary.⁴ For a definitive diagnosis, PCR of a serum or eschar sample is useful. If appropriate treatment, that is, tetracyclines or macrolides,⁵ is not administered early, the incidence of severe complications such as disseminated intravascular coagulation generally becomes high,⁴ although the outcome in immunocompromised patients such as those on dialysis and on immunosuppressants is not well understood. In conclusion, when fever develops after farming or gardening in late autumn in *Orientia tsutsugamushi*-rich regions, careful examination for a possible eschar is of significance for early diagnosis and treatment.

Learning points

- ▶ Scrub typhus is transmitted by the bite of a tick larva, called chigger, and after an incubation period of 7–10 days, symptoms such as headache, myalgia and rash are observed besides fever and chills. While the accompanying symptoms other than fever and chills may be unremarkable, they are often fatal.
- ▶ In scrub typhus, approximately 80% of patients have an eschar, which is of high diagnostic value, but patients themselves do not often notice the presence of the eschar, which may be found in obscure sites such as the pubis, axilla or lower legs; therefore, careful physical examination is necessary.
- ▶ When fever develops after farming or gardening in late autumn in *Orientia tsutsugamushi*-rich regions, careful examination for a possible eschar is of significance for early diagnosis and treatment.



Figure 1 Detailed physical examination revealed an eschar of 5 mm diameter on the medial side of the right lower leg.

Contributors TA contributed to write manuscript, patient care and discussion. KI, NT and IM also contributed to discussion and patient care.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.



© BMJ Publishing Group Limited 2023. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Akaike T, Ishizuka K, Tominaga N, et al. *BMJ Case Rep* 2023;**16**:e255404. doi:10.1136/bcr-2023-255404

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Kosuke Ishizuka <http://orcid.org/0000-0003-4313-6592>

REFERENCES

- 1 Bonell A, Lubell Y, Newton PN, *et al*. Estimating the burden of scrub typhus: a systematic review. *PLoS Negl Trop Dis* 2017;11:e0005838.
- 2 Sando E, Suzuki M, Katoh S, *et al*. Distinguishing Japanese spotted fever and scrub typhus, central Japan, 2004–2015. *Emerg Infect Dis* 2018;24:1633–41.
- 3 Kinoshita H, Arima Y, Shigematsu M, *et al*. Descriptive epidemiology of rickettsial infections in Japan: scrub typhus and Japanese spotted fever, 2007–2016. *Int J Infect Dis* 2021;105:560–6.
- 4 Shikino K, Ohira Y, Ikusaka M. Scrub typhus (tsutsugamushi disease) presenting as fever with an eschar. *J Gen Intern Med* 2016;31:582.
- 5 El Sayed I, Liu Q, Wee I, *et al*. Antibiotics for treating scrub typhus. *Cochrane Database Syst Rev* 2018;9:CD002150.

Copyright 2023 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow