[Images in...](#)

The best treatment can be no treatment: retained retro-orbital air gun pellet following attempted suicide

Hugh Jewsbury,¹ Des O'Duffy²

¹Department of Ophthalmology, HM Stanley Hospital, St Asaph, UK;

²Department of Ophthalmology, Royal Gwent Hospital, Newport, UK

Correspondence to Hugh Jewsbury, hughjewsbury@hotmail.com

DESCRIPTION

A 35-year-old man presented after shooting himself in the right orbit with an air rifle following an attempted suicide. His visual acuity was 6/5 in the right eye and 6/6 in the left. An entry wound was noted on the medial aspect of the upper lid. The anterior segment was otherwise unremarkable. Examination of the posterior segment revealed

no significant abnormalities. There was no restriction of ocular motility. X-rays demonstrated a retro-orbital metallic foreign body (figure 1). This was shown to lie above the lateral rectus and behind the globe on a subsequent CT scan (figure 2). He was treated with broad-spectrum antibiotics and the pellet was not removed as the risk of surgery outweighed the benefit.

Figure 1 Plain x-ray of facial bones showing retained air rifle pellet.

Figure 2 CT revealed the precise location above the right lateral rectus muscle and behind the globe.

The retained pellet was composed of lead. There are no reported cases of systemic lead toxicity following orbital retention but this remains a theoretical risk.¹ Most metals are inert but certain metals pose specific problems. Retention of ferromagnetic metals will impede future MRI, while retention of iron is associated with siderosis and copper with purulent orbital inflammation.¹ Unless there is a specific indication relating to the type of metal retained, removal of retro-orbital, inorganic foreign bodies should only be considered in the presence of adverse effects from retention. These indications include orbital inflammation, infection, optic nerve compromise or restricted ocular motility.^{1–3}

This is a very unusual and fortuitous presentation of an ocular air gun injury and demonstrates that conservative management in such circumstances can be most effective.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Ho VH**, Wilson MW, Fleming JC, et al. Retained intraorbital metallic foreign bodies. *Ophthalm Plast Reconstr Surg* 2004;**20**:232–6.
2. **Fulcher TP**, McNab AA, Sullivan TJ. Clinical features and management of intraorbital foreign bodies. *Ophthalmology* 2002;**109**:494–500.
3. **Finkelstein M**, Legmann A, Rubin PA. Projectile metallic foreign bodies in the orbit: a retrospective study of epidemiologic factors, management, and outcomes. *Ophthalmology* 1997;**104**:96–103.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Jewsbury H, O'Duffy D. The best treatment can be no treatment: retained retro-orbital air gun pellet following attempted suicide. *BMJ Case Reports* 2011;10.1136/bcr.02.2011.3826, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow