Images in...

320-slice CT and myocardial bridge

Antonietta Evangelista, Francesco Pelliccia, Alessio Arrivi, Carlo Gaudio

Department of Heart and Great Vessels, University Sapienza, Rome, Italy

Correspondence to Dr Alessio Arrivi, alessio.arrivi@libero.it

DESCRIPTION

A 29-year-old professional athlete was referred for cardiac evaluation after a syncope occurring during strenuous exercise. Twelve-lead ECG showed Brugada-like pattern (figure 1), but both flecainide and genetic tests were negative. ECG Holter and echocardiogram were normal. ECG stress test showed no significant ST-segment changes. In order to rule out coronary artery disease or anomalies, 320-slice multi detector CT (Aquilion One; Toshiba Medical Systems Inc, Tokyo, Japan) was done with settings at 100 mV and 320 mA (heart rate was 50 beats/min). Eighty millilitre of contrast agent (Ultravist 370; Bayer Schering Pharma AG, Berlin, Germany) were injected intravenously at a rate of 5.5 ml/s, followed by 60 ml of saline at the same rate. The total radiation dose

was only 2.4 mSv. The scan showed two typical myocardial bridges of the left anterior descending artery, with intramural tracts of 10 mm in the mid-segment and of 15 mm in the distal segment (figure 2, upper and lower panels). No evidence of stenosis of coronary arteries was found.

CT can rule out coronary artery disease in very low-risk subjects and detect myocardial bridging.¹ With the introduction of 320-slice multi-slice CT, scanning of the entire heart has become possible in a single heart beat with a very low dose of radiation.² Thus, this novel technique is particularly suited for cases (ie, such as young professional athletes) in whom is essential to rule out any possible coronary artery disease or anomaly avoiding more invasive procedures.

Figure 1 Twelve-lead ECG showing Brugada-like pattern.

Figure 2 The 320-slice scans showed two typical myocardial bridges of the left anterior descending artery, with intramural tracts of 10 mm in the mid-segment and of 15 mm in the distal segment (arrows).

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Faletta F**, Auricchio A, Moccetti T, *et al*. "Dynamic imaging" (systolic compression) of myocardial bridge visualised by electronic beam computed tomography. *Heart* 2007;**93**:1135.
2. **de Graaf FR**, Schuijff JD, Delgado V, *et al*. Clinical application of CT coronary angiography: state of the art. *Heart Lung Circ* 2010;**19**:107–16.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Evangelista A, Pelliccia F, Arrivi A, Gaudio C. 320-slice CT and myocardial bridge. *BMJ Case Reports* 2011;10.1136/bcr.05.2011.4200, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow