Images in...

Gastrointestinal bleeding caused by gastritis cystica polyposa

Kenji Tominaga,¹ Satoshi Nimura,² Kei Takahashi,³ Iruru Maetani¹¹Department of Internal Medicine, Division of Gastroenterology, Toho University Ohashi Medical Center, Tokyo, Japan;²Department of Pathology, Faculty of Medicine, Fukuoka University, Fukuoka, Japan;³Department of Surgical Pathology, Toho University Ohashi Medical Center, Tokyo, Japan

Correspondence to Dr Kenji Tominaga, ktominaga@oha.toho-u.ac.jp

DESCRIPTION

A man in his 50s presented with a 2-day history of melaena. He had undergone a Billroth II (B-II) gastrectomy for a benign duodenal ulcer 32 years ago. He had a history of heavy alcohol intake and smoked 20 cigarettes per day. Laboratory testing revealed anaemia. Oesophagogastroduodenoscopy revealed a small amount of blood in the gastric remnant and a polypoid lesion at the B-II gastroenterostomy site. Bleeding at the polypoid lesion could be seen on contact with the endoscope. However, no treatment was needed during the oesophagogastroduodenoscopy because of spontaneous haemostasis. A biopsy specimen from the lesion demonstrated foveolar epithelial hyperplasia and marked cystic dilatation of pyloric glands (figure 1). The patient was transfused with four units of packed red blood cells. He ceased alcohol intake and cigarette smoking and did well without further blood transfusion. No bleeding was seen at 46 days (figure 2) after the initial oesophagogastroduodenoscopy and endoscopic ultrasonography revealed multiple cystic lesions mainly in a thickened submucosal layer. Based on these results, the patient was diagnosed with gastrointestinal bleeding caused by gastritis cystica polyposa (GCP). GCP is usually

observed more than 10 years after B-II gastrectomy.¹ The disease is characterised by gastrointestinal bleeding and coexistence with gastric remnant carcinoma. In cases with massive bleeding and/or suspicion of malignancy, endoscopic haemostasis or surgery are the treatments of choice. GCP is considered to be a precancerous lesion, therefore follow-up of patients with GCP is essential to detect gastric remnant carcinoma including gastric carcinoma arising at the site of GCP.^{2 3}

Competing interests None.

Patient consent Obtained.

REFERENCES

1. Littler ER, Gleibermann E. Gastritis cystica polyposa. (Gastric mucosal prolapse at gastroenterostomy site, with cystic and infiltrative epithelial hyperplasia). *Cancer* 1972;**29**:205–9.
2. Kaizaki Y, Hosokawa O, Sakurai S, et al. Epstein-Barr virus-associated gastric carcinoma in the remnant stomach: de novo and metachronous gastric remnant carcinoma. *J Gastroenterol* 2005;**40**:570–7.
3. Matsushita M, Mori S, Tahashi Y, et al. Gastritis cystica polyposa in the operated stomach and heterotopic submucosal cysts in the unoperated stomach. *Gastrointest Endosc* 2010;**71**:1100–1.

Figure 1 A biopsy specimen from the lesion demonstrates elongated, tortuous foveolae and hyperplastic and cystically dilated underlying pyloric glands. The lamina propria is slightly oedematous and contains a mild chronic inflammatory cellular infiltrate (H&E, original magnification x20).

Figure 2 Endoscopic retroflex view of the Billroth II gastroenterostomy site at 46 days after the initial oesophagogastroduodenoscopy. A polypoid lesion (ie, gastritis cystica polyposa) can be seen with no bleeding.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Tominaga K, Nimura S, Takahashi K, Maetani I. Gastrointestinal bleeding caused by gastritis cystica polyposa. *BMJ Case Reports* 2011; 10.1136/bcr.08.2011.4617, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow