Images in...

Pectus excavatum

Umar Imran Hamid, Harry Parissis

Department of Cardiothoracic Surgery, Royal Victoria Hospital, Belfast, UK

Correspondence to Umar Imran Hamid, umar79@hotmail.com

DESCRIPTION

Posterior depression of the sternum and costal cartilages produces the characteristic findings of pectus excavatum: funnel chest, or *trichterbrust*. Pectus excavatum is present at birth or within the first year of life in the majority of affected children. It occurs more frequently in boys than girls, by almost a 4:1 ratio. Although the sternal depression appears to be caused by an overgrowth of costal cartilages, the aetiology of pectus deformities is unknown. Deformity of the chest wall led many authors to attribute the symptomatic improvement in patients after surgery. Posterior displacement of the sternum can produce a deformity of the heart, particularly anterior indentation of the right ventricle. The debate is ongoing among clinicians regarding improvement in cardiopulmonary function following repair.¹ Correction of pectus deformity ranges from minimally invasive techniques like Nuss procedure² to open repairs.³ Complications of pectus excavatum repair are few and relatively unimportant except for major recurrence.⁴

Of clinical significance is the presence of various cardiac abnormalities ranging from valvular dysfunction to tetralogy of fallot. The images are typical depiction of a pectus deformity with posterior displacement of the body of the sternum.

Competing interests None.

Patient consent Not obtained.

REFERENCES

1. **Malek MH**, Fonkalsrud EW, Cooper CB. Ventilatory and cardiovascular responses to exercise in patients with pectus excavatum. *Chest* 2003;**124**:870–82.
2. **Nuss D**, Kelly RE Jr, Croitoru DP, *et al*. A 10-year review of a minimally invasive technique for the correction of pectus excavatum. *J Pediatr Surg* 1998;**33**:545–52.
3. **Fonkalsrud EW**, Beanes S, Hebra A, *et al*. Comparison of minimally invasive and modified Ravitch pectus excavatum repair. *J Pediatr Surg* 2002;**37**:413–17.
4. **Shamberger RC**, Welch KJ. Surgical repair of pectus excavatum. *J Pediatr Surg* 1988;**23**:615–22.

Figure 1 (A,B) CT images indicating typical posterior displacement of sternum in pectus excavatum.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Hamid UI, Parissis H. Pectus excavatum. *BMJ Case Reports* 2011;10.1136/bcr.12.2010.3581, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow