Images in...

Juvenile oral lichen planus: a clinical rarity

T R Chaitra,¹ Ravishankar Lingesh Telgi,² Amit Kishor,¹ Adwait Uday Kulkarni³¹Department of Pediatric Dentistry, Kothiwal Dental College and Research Centre, Moradabad, Uttar Pradesh, India²Department of Public Health Dentistry, Kothiwal Dental College and Research Centre, Moradabad, Uttar Pradesh, India³Oral and Maxillofacial Surgery Department, Sinhgad Dental College, Pune, Maharashtra, India

Correspondence to Dr Adwait Uday Kulkarni, dradwaitkulkarni@gmail.com

DESCRIPTION

A 9-year-old girl reported to the department with the chief complaint of burning sensation in the buccal mucosa of right and left side. Burning sensation was there only during consumption of spicy food. History and general physical examination revealed no relevant findings. On oral examination, it was evidently seen that there was white reticular lines and dark red and white patches on both the left and right buccal mucosa (figures 1 and 2). On palpation, there was mild pain. Provisional diagnosis of lichen planus (LP) was made by its presentation. Erosive LP, reticular LP, allergic mucositis, lichenoid reaction and thermal burn were given as differential diagnosis. Punch biopsy was performed and the histopathological findings revealed the presence of stratified squamous epithelium with inflamed connective tissue stroma and few superficial areas of erosion (figure 3). Final diagnosis of erosive LP was made. Topical corticosteroid (triamcinolone acetonide 0.1%) and analgesic (2% lidocain gel) were prescribed. A marked reduction in symptom was reported after one week. LP was first described in the literature by Erasmus Wilson in 1869, as predominately a disease of the middle aged or older. There is limited literature available reporting the occurrences of oral LP in children. The aetiology remains uncertain but many factors have been implicated such as genetic predisposition, infective agents, systemic diseases, graft versus host disease, drug reactions, hypersensitivity to dental materials and vitamin deficiencies. Also, it has been documented as a complication of

Figure 2 Shows right buccal mucosa.

hepatitis B vaccinations¹ and has autoimmune basis.² The exact incidence of paediatric LP is unknown. Several retrospective reviews, have estimated that only 1–16% of LP

Figure 1 Shows left buccal mucosa.

Figure 3 H&E staining of biopsy.

patients are younger than 15 years old, has rarely been documented in the medical/dental literature.² Regarding the clinical features, some authors suggest that they are similar to adult LP, but some others suggest that in children it was often atypical with a linear pattern.³ Of note, tacrolimus ointment, topical tretinoin and topical cyclosporine are used with success in some cases for treatment. Periodic follow-up is required in all oral lichen planus in children, as malignant transformation is seen in small percentage of cases on follow-up.²

Learning points

- ▶ Oral lichen planus is rare in children.
- ▶ If it occurs in this age group, pedodontists should be watchful to first diagnose and give appropriate treatment.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Patil S**, Yeoman CM, Murphy R. Oral lichen planus in childhood: a report of three cases. *IJPD* 2005;**15**:118–22.
2. **Woo VL**, Manchanda-Gera A, Park DS, *et al*. Juvenile oral lichen planus: a report of 2 cases. *Pediatr Dent* 2007;**29**:525–30.
3. **Alam F**, Hamburger J. Oral mucosal lichen planus in children. *IJPD* 2001;**11**:209–14.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Chaitra TR, Telgi RL, Kishor A, Kulkarni AU. Juvenile oral lichen planus: a clinical rarity. *BMJ Case Reports* 2012;10.1136/bcr-2012-006192, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow