

Images in...

Plummer-Vinson syndrome

Sanat Phatak, Neelam Redkar, Meenakshi Amit Patil, Ajay Kuwar

Department of Medicine, Seth G S Medical College and KEM Hospital, Mumbai, Maharashtra, India

Correspondence to Dr Meenakshi Amit Patil, meenakshi.patil90@gmail.com, meenakshi.patil90@yahoo.com

DESCRIPTION

A 27-year-old woman presented to the medicine outpatient clinic with easy fatiguability and breathlessness on exertion since 2 months. She also complained of sore oral mucosa and was unable to swallow large morsels of food. Physical examination revealed angular cheilitis and pigmentation of the oral mucosa, severe pallor, koilonychia (figure 1), large volume pulse and a haemic murmur in the left parasternal area. Investigations revealed hypochromic microcytic anaemia with haemoglobin of 5.6 g%. Serum iron studies, showed an increased total iron binding capacity (TIBC) of 512 $\mu\text{g/l}$ (250–370 $\mu\text{g/l}$) and low transferrin saturation 10% (15–20%) suggesting iron-deficiency anaemia. Barium swallow and upper gastrointestinal endoscopy was done to assess the cause of dysphagia. Barium swallow showed upper oesophageal webs (figure 2). On upper gastrointestinal endoscopy, post cricoid oesophageal web was seen and the endoscope could not be passed across the cricoid level (figure 3). Fulfilling the classical triad of dysphagia, iron-deficiency anaemia and upper oesophageal webs, Plummer-Vinson syndrome (or Paterson-Brown Kelly syndrome, sideropenic dysphagia) was diagnosed. In subsequent endoscopy, the webs were dilated with considerable resultant improvement in her dysphagia. Obvious ulcerated heterotopic gastric mucosa

patches were ruled out. Iron supplementation was initiated and the patient was counselled regarding the need for surveillance for the development of an upper gastrointestinal malignancy.

The pathogenesis of Plummer-Vinson syndrome remains speculative. Iron deficiency has been implicated in the pathogenesis of oesophageal webs and dysphagia in predisposed individuals. The depletion of iron-dependent oxidative enzymes may produce myasthenic changes in muscles involved in the swallowing mechanism, atrophy of the oesophageal mucosa, and formation of webs as epithelial complications. The improvement in dysphagia after iron therapy provides evidence for an association between iron deficiency and postcricoid dysphagia.¹ Moreover, the decline in Plummer-Vinson syndrome seems to parallel an improvement in nutritional status, including iron supplementation. Post-cricoid web or Plummer-Vinson syndrome has been identified as a risk factor for the development of upper gastrointestinal tract malignancy.²



Figure 1 Severe koilonychia.



Figure 2 Upper oesophageal webs seen on barium swallow.




Figure 3 Upper gastrointestinal endoscopy showing oesophageal web.

Learning points

- ▶ Plummer-Vinson syndrome classically presents as a triad of iron-deficiency anaemia, postcricoid dysphagia and upper esophageal webs.
- ▶ Treatment of the iron deficiency frequently relieves dysphagia though some patients may need endoscopic web dilatation.
- ▶ Patients must be screened regularly for any development of upper gastrointestinal malignancy.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Dantas RO**, Villanova MG. Esophageal motility impairment in Plummer-Vinson syndrome. Correction by iron treatment. *Dig Dis Sci* 1993;**38**:968–71.
2. **Chisholm M**. The association between webs, iron and post-cricoid carcinoma. *Postgrad Med J* 1974;**50**:215–19.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
 BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Phatak S, Redkar N, Patil MA, Kuwar A. Plummer-Vinson syndrome. *BMJ Case Reports* 2012;10.1136/bcr-2012-006403, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow