Images in...

Pyopneumopericardium

Smitha Bhat, Jayaprakash Alva, Shishir Duble, Muhamed Ali

Department of Medicine, Father Muller Medical College, Mangalore, Karnataka, India

Correspondence to Dr Smitha Bhat, doctorsmitha@yahoo.co.in

DESCRIPTION

An elderly woman with asthma presented with symptoms of dyspepsia. Gastroscopy showed a hiatus hernia with large gastric ulcer. She was treated with proton pump inhibitors and improved symptomatically. Four weeks later, she developed sudden-onset breathlessness and wheeze. The patient was afebrile. Initial examination revealed tachycardia, tachypnoea and severe bronchospasm. No evident focus of sepsis was found at this point, and patient improved after treatment with bronchodilators. A day later, however, patient's breathlessness worsened and a chest x-ray revealed a pneumopericardium. Echocardiogram showed the presence of pneumopericardium, with no evidence of tamponade (figure 1). This was further confirmed by CT scan of the chest, which revealed pneumopericardium. However,

pericardiocentesis revealed the presence of pus in addition to air. Pus grew *Enterococcus faecalis*. An endoscopy was performed, which revealed an ulcer in the hiatus hernia perforating into the pericardial space. The poor general condition of the patient precluded surgical intervention and she passed away in spite of broad-spectrum antibiotics and drainage of pus in the pericardial sac.

Pyopneumopericardium is a serious condition that has a death rate of almost 100%.¹ Surgery is the treatment of choice for pneumopericardium due to gastropericardial fistula regardless of the site of the fistula.²

Learning points

- Early pericardiocentesis is essential to diagnose pyopneumopericardium even when imaging reveals only pneumopericardium.
- Surgery must be considered as the treatment of choice even in patients with poor preoperative physical status.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Cummings RG**, Wesly RLR, Adams DH, *et al* Pneumopericardium resulting in cardiac tamponade. *Ann Thorac Surg* 1984;**37**:511–18.
2. **Salling N**, Falensteen AM, Larsen LG. Non-traumatic perforation of gastric ulcer in a hiatal hernia to the pericardium. *Acta Med Scand* 1983;**213**:225–6.

Figure 1 Pyopneumopericardium.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Bhat S, Alva J, Doble S, Ali M. Pyopneumopericardium. *BMJ Case Reports* 2012;10.1136/bcr-2012-006411, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow