Images in...

Polysling skin pressure necrosis—a complication of shoulder immobilisation

Sarkhell Radha, Kalpesh R Vaghela, Sujith Konan, Warwick Radford

Department of Trauma and Orthopaedic Surgery, Chelsea & Westminster Hospital, London, UK

Correspondence to Dr Kalpesh R Vaghela, kalpeshrvaghela@gmail.com

DESCRIPTION

A 65-year-old man sustained a closed minimally displaced fracture of the right proximal humerus following a low-velocity road traffic accident. The alignment of the humeral head was satisfactory and the patient was treated conservatively with a polysling shoulder immobiliser (TalarMade, Universal Multi Arm Sling). He was reviewed in fracture clinic 2 weeks postinjury where the position and alignment of the fracture fragments remained satisfactory on check radiographs. Subsequent review at 4 weeks revealed a 10×2 cm² grade 3 pressure sore and surrounding erythema caused by the polysling strap (figure 1). The finding was incidental and he was completely asymptomatic to the effect of the strap on his neck. Following this, the strap was adjusted and additional padding was used to relieve pressure. The wound was managed by a district

Figure 1 Grade 3 pressure sore caused by the polysling strap.

Figure 2 Resolution of the pressure sore at 3-week follow-up.

nurse with dressings which saw resolution of the pressure sore at 3 weeks (figure 2).

Polyslins are nylon slings that are commonly used to immobilise the shoulder and support the upper limb. Patients and healthcare professionals should be alerted to this potential complication and advised to assess the integrity of the skin and regularly inspect pressure areas. In this case, the patient was a healthy male with no additional risk factors for development of pressure sores. Risk assessments such as the Braden Score¹ can be used to identify high-risk patients. Treatment includes relief of pressure, dressings, debridement and antibiotics in cases with superimposed infection but as always prevention is better than cure.²

Learning points

- ▶ Immobilisation of fractures and dislocations using slings and casts can lead to the preventable complication of pressure sores.
- ▶ Risk assessment tools such as the Braden Score can be used to identify patients who are at risk of developing pressure sores.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Bergstrom N**, Braden BJ. The Braden scale for predicting pressure sore risk. *Nursing Res* 1987;**36**:205–10.
2. Pressure ulcers: The management of pressure ulcers in primary and secondary care. NICE clinical guideline 2005.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Radha S, Vaghela KR, Konan S, Radford W. Polysling skin pressure necrosis—a complication of shoulder immobilisation. *BMJ Case Reports* 2012;10.1136/bcr-2012-006671, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow