Images in...

Multiple cerebral hydatid cysts: have the previous operations contributed to their formation?

Osama S M Amin

Department of Neurology, Sulaimaniya General Teaching Hospital, Sulaimaniya City, Iraq

Correspondence to Dr Osama S M Amin, dr.osama.amin@gmail.com

DESCRIPTION

A 32-year-old Iraqi Kurdish woman was brought to A&E after developing recurrent generalised tonic-clonic seizures. During the past 4 years, she underwent two craniectomies in Iran for cerebral hydatid cystic disease. Oral albendazole was prescribed for 3 months only, prior to the first surgical operation. The first operation and its discharging notes, including a histopathological report, revealed that she had a solitary hydatid cyst (of *Echinococcus granulosus*) of the right parietal hemispheric area. The neurosurgeon used short courses of glucocorticoids before and after both craniectomies. Her brain MRI examination is shown in figures 1 and 2.

Brain involvement occurs in 1–2% of all *E granulosus* infections. The cysts are usually located supratentorially.

Figure 1 Axial T2-weighted brain MRI of the patient. There are multiple cysts of variable sizes and shapes at the right parieto-occipital lobes. They are surrounded by perilesional oedema (black arrow). Three sites of the previous craniectomies can be seen (yellow arrows). Several cysts demonstrate internal septations, reflecting the presence of 'daughter cysts'. Flow void signals (red arrow) can be observed at the periphery of some cysts indicating calcification. The contents of the cysts have a hyperintense (cerebrospinal fluid-like) signal. The patient was given oral albendazole tablets for 3 months prior to the first operation only; since then, this medication had never been prescribed. Glucocorticoids were administered during the perioperative periods of her two craniectomies.

Children and young adults are the usual targets. Multiple intracerebral hydatid cysts, whether primary or secondary, are uncommon. In the series of Onal *et al*,¹ 4 of their 45 patients with solitary intracranial hydatid cysts developed recurrence after neurosurgical removal; all of them followed intraoperative rupture of a primary solitary supratentorial hydatid cyst.

Krajewski and Stelmasiak² studied 12 patients with cerebral hydatid cystic disease who underwent brain surgery and they followed up them for 5 years. They found that recurrence of multiple cysts occurred in one case and another patient was reoperated on twice for recurrent cysts after an operation in another centre.

It should be noted that despite all the advances in imaging techniques and therapeutic methods, central nervous system hydatidosis remains difficult to cure and patient outcomes are not satisfactory due to the high incidence of recurrence.³

Learning points

- ▶ Multiple intracerebral hydatid cysts are rare.
- ▶ The formation of multiple cerebral hydatid cysts after neurosurgical removal of a primary solitary cyst reflects an intraoperative rupture of the latter with secondary development of these hydatid cysts.
- ▶ Central nervous system hydatidosis remains difficult to cure and patient outcomes are not satisfactory due to the high incidence of recurrence.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. Onal C, Unal F, Barlas O, *et al*. Long-term follow-up and results of thirty pediatric intracranial hydatid cysts: half a century of experience in the Department of Neurosurgery of the School of Medicine at the University of Istanbul (1952–2001). *Pediatr Neurosurg* 2001;**35**:72–81.
2. Krajewski R, Stelmasiak Z. Cerebral hydatid cysts in children. *Childs Nerv Syst* 1991;**7**:154–5.
3. Limaiem F, Bellil S, Bellil K, *et al*. Primary hydatidosis of the central nervous system: a retrospective study of 39 Tunisian cases. *Clin Neurol Neurosurg* 2010;**112**:23–8.

Figure 2 Coronal T2-fluid attenuation inversion recovery (FLAIR) brain MRI film of the patient. There are right-sided parietal cysts (with internal septae of daughter cysts) surrounded by vasogenic oedema of the white matter. The hyperintense signal content of the cysts (on the T2-weighted film) was 'suppressed' on this FLAIR sequence, indicating that these cysts contain a cerebrospinal fluid-like watery fluid. The patient was admitted to the intensive care unit for status epilepticus management; no kind of any neurosurgical intervention was done. She died on day 2.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit

<http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Amin OSM. Multiple cerebral hydatid cysts: have the previous operations contributed to their formation?. *BMJ Case Reports* 2012;10.1136/bcr-2012-007240, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow