Quadriceps fat-pad impingement syndrome: MRI findings

Ahmet Bas, Onur Tutar, Inanc Yanik, Cesur Samanci

Department of Radiology,
Istanbul University Cerrahpasa
Medical Faculty,
Istanbul, Turkey

Correspondence to

Dr Ahmet Bas,
dr.ahmetbas@hotmail.com

DESCRIPTION

A 32-year-old man was referred to a radiology department for intermittent anterior knee pain and swelling for approximately 6 months. The patient denied any history of major trauma, fracture or infection. MRI was performed to diagnose the cause of knee pain. The axial view of MRI revealed increased patellofemoral joint effusion and the sagittal view of T1-weighted, and T2-weighted MRIs showed an oedema, increased intra-articular fluid and thickening of suprapatellar fat pad

(figures 1 and 2). Also, oedema of quadriceps fat impingement is visible at both coronal and axial fat sat T2-weighted sections.

Knee pain is a very common clinical complaint. Generalised pain in the front of knee has many causes. Numerous traumatic and non-traumatic processes may be encountered, many occurring relative to the anterior compartment, including fat-pad impingement syndromes. Three normal anterior knee fat pads are present at the knee which separates joint capsule and synovium; the

Figure 1 The axial T2 fat sat image demonstrates increased intra-articular fluid.

Figure 3 Quadriceps fat pad (arrow) is hyperintense on the sagittal T2 fat sat image (A) and hypointense on sagittal T1 image (B).

Figure 2 The axial T2 fat sat image demonstrates increased signal and thickening (arrow) within fat interposed between the quadriceps tendon and the suprapatellar bursa.

To cite: Bas A, Tutar O, Yanik I, et al. *BMJ Case Reports* Published online: 10 December 2012
doi:10.1136/bcr-2012-007643

Figure 4 The coronal T2 fat sat image demonstrates increased signal within the quadriceps fat pad and mild effusion surrounding (patella=green arrow, quadriceps fat pad=yellow arrow, effusion=red arrow).

quadriceps (anterior suprapatellar), the prefemoral (posterior suprapatellar or supratrochlear) and Hoffa (infrapatellar) fat pads.^{1 2} The quadriceps fat pad is the normal fat pad between the suprapatellar recess posteriorly and quadriceps tendon anteriorly. The term 'quadriceps fat-pad impingement' has been used to

describe an inflammatory process within the anterior suprapatellar fat, manifested on MRI as high T2 signal, low T1 signal and mass effect on the quadriceps tendon³ (figures 3 and 4).

Learning points

- ▶ Several entities with anterior knee pain that involve peripatellar fat may be encountered when evaluating MR of the knee. Of these, one of the most common is quadriceps/suprapatellar fat-pad impingement syndrome which may be related to chronic microtrauma.
- ▶ The term 'quadriceps fat-pad impingement' has been used to describe an inflammatory process within the anterior suprapatellar fat, manifested on MRI as high T2 signal, low T1 signal and mass effect on the quadriceps tendon.

Competing interests None.

Patient consent Obtained.

REFERENCES

- 1 Jacobson JA, Lenchik L, Ruhoy MK, *et al.* MR imaging of the infrapatellar fat pad of Hoffa. *Radiographics* 1997;17:675–91.
- 2 Schweitzer ME, Falk A, Pathria M, *et al.* MR imaging of the knee: can changes in the intracapsular fat pads be used as a sign of synovial proliferation in the presence of an effusion". *AJR* 1993;160:823–6.
- 3 Shabshin N, Schweitzer ME, Morrison WB. Quadriceps fat pad edema significance on magnetic resonance images of the knee. *Skeletal Radiol* 2006;35:269–74.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow