Images in...

Enterocolitis with multiple ulcers of ileum and right colon in a patient with leukaemia attributed to cytomegalovirus

Iordanis N Papadopoulos, Ioanna Konstantiadou, Evaggelia Papantoni

Fourth Surgery Department, National & Kapodistrian University of Athens, Attikon University General Hospital, Athens, Greece

Correspondence to Professor Iordanis N Papadopoulos, ipapado@med.uoa.gr

DESCRIPTION

A 76-year-old woman with a history of myelogenous leukaemia was admitted with haematochezia fever (38.3°C) and slight abdominal pain. Haemoglobin was 8 g/l, WBC $8.2 \times 10^9/l$, platelets were $55 \times 10^9/l$ and partial thromboplastin time was 23 s. Gastroscopy was normal, colonoscopy did not reveal the source of haemorrhage, but a CT-angiography indicated the source at the ileo-caecal region.

Non-surgical therapy including 8 units of blood and 8 units of fresh frozen plasma was performed in 3 days, but haemorrhage continued, (HB 6 g/l).

A right hemicolectomy was performed including en-block resection of 40 cm of the terminal ileum. Intraluminal absence of blood and focal thickening around the ulcers guided the extent of resection. The presence of multiple ulcers was confirmed before closing the abdomen, figures 1 and 2. Histology revealed no malignant infiltration, but inflammation with intracellular inclusion bodies and serology consisting of IgG anti-cytomegalovirus (CMV), and IgM anti-CMV indicated the CMV as the aetiology of the ulcers.

Postoperatively, the haemorrhage did not recur but the patient developed a low-output entero-cutaneous fistula (3rd week). She initially showed improvement by total parenteral nutrition, antibiotics and ganciclovir.

Nevertheless, the patient's haematological status remained poor and she died 70 days after surgery in sepsis due to pneumonia.

CMV^{1 2} is a herpes virus with a propensity to affect immunocompromised patients as transplant recipients, patients with AIDS, haematological malignancies and inflammatory bowel disease. Pneumonia, gastrointestinal inflammation, ulcerations and haemorrhage, hepatitis and

Learning points

- ▶ The multiplicity and wide spread of ileal and colonic ulcers make the assessment of the extent of surgical resection difficult.
- ▶ Surgery should be reserved for life-threatening complications as haemorrhage and perforation.
- ▶ Long-term prognosis is largely determined by the underlying haematologic disease.

Figure 1 Segment of the resected terminal ileum with three distinct ulcers. Each ulcer is indicated by four white arrows. The crater of each ulcer is surrounded by an opaque oedematous ridge and the convergence of circular plicae.

Figure 2 Segment of the resected caecum and ascending colon showing two distinct ulcers of colitis. Each ulcer is indicated by four white arrows. The crater of each ulcer is surrounded by an opaque oedematous ridge and the convergence of the intestinal plicae. The multiple mucosal red spots represent inflammatory foci and mucosal erosions. The red arrows indicate the opened ileocaecal valve.

encephalitis are the most common complications with high mortality. Antiviral therapy (ganciclovir, valganciclovir) should be considered.³

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Boeckh M**, Geballe AP. Cytomegalovirus: pathogen, paradigm, and puzzle. *J Clin Invest* 2011;**121**:1673–80.
2. **Lawlor G**, Moss AC. Cytomegalovirus in inflammatory bowel disease: pathogen or innocent bystander? *Inflamm Bowel Dis* 2010;**16**:1620–7.
3. **O'Brien S**, Ravandi F, Riehl T, *et al.* Valganciclovir prevents cytomegalovirus reactivation in patients receiving alemtuzumab-based therapy. *Blood* 2008; **111**:1816–19.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit

<http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Papadopoulos IN, Konstantiadou I, Papantoni E. Enterocolitis with multiple ulcers of ileum and right colon in a patient with leukaemia attributed to cytomegalovirus. *BMJ Case Reports* 2012;10.1136/bcr.01.2012.5651, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow