Images in...

Gallstone ileus

Miguel F Carrascosa,¹ Mónica D Riego-Martín,² José-Ramón Salcines Caviedes,³ Pablo González Gutiérrez⁴

¹Department of Internal Medicine, Hospital of Laredo, Laredo, Cantabria, Spain;

²Department of Radiology, Hospital of Laredo, Laredo, Cantabria, Spain;

³Department of Digestive Diseases, Hospital of Laredo, Laredo, Cantabria, Spain;

⁴Department of Radiology Service, Hospital of Laredo, Laredo, Cantabria, Spain

Correspondence to Dr Miguel F Carrascosa, miguel.carrascosa@scsalud.es

DESCRIPTION

The diagnosis of gallstone ileus entails a clinical challenge, especially in older patients in whom it may be easily overlooked.¹ Herein, we report on a patient suffering from this condition who successfully recovered after surgery. A 65-year-old man presented to our emergency department with nausea, recurrent vomiting and colicky abdominal pain of 5 days' duration. His medical history was relevant only for hypertension. Physical examination revealed a non-tender abdomen with active bowel sounds. Laboratory tests showed mild leukocytosis and severe, acute renal impairment. Abdominal radiograph demonstrated the presence of a round hyperdense mass with calcified margins in the right lower quadrant as well as enlarged loops of small bowel (figure 1). CT scan of the abdomen showed pneumobilia, a calcified intraluminal mass impacted in the terminal ileum and dilated upstream

loops of small intestine (figure 2). These findings were diagnostic for gallstone ileus. On the same admission day, the patient was transferred to the operating room where a simple enterotomy and removal of the obstructing gallstone (5.2×3.6 cm) were performed. He made an uneventful recovery and was discharged 8 days after surgery. Gallstone ileus is an unusual complication of cholelithiasis that particularly affects older individuals and continues to be associated with high morbidity and mortality. It is caused by intestinal impaction of one or more gallstones after being passed through a biliary-enteric fistula. Most ectopic gallstones impact the ileum, which is the narrowest portion of the intestine. While gallstone ileus is responsible for only 1–4% of all cases of mechanical bowel obstruction,² it causes up to 25% of non-strangulated small bowel obstruction in patients over age 65.³ Thus, gallstone ileus should always be high in the differential diagnosis when assessing an aged person with intestinal obstruction. The disorder, however, is often not considered since patients frequently deny a history of biliary disease. Strong clinical

Figure 1 Plain abdominal radiograph of a 65-year-old man showing signs of small-bowel mechanical obstruction caused by an ectopic gallstone (arrow).

Figure 2 CT scan of the abdomen revealing pneumobilia, a gallstone in the distal ileum (arrow) and small-bowel dilatation.

suspicion, timely use of combined diagnostic imaging modalities (mainly, abdominal plain film and CT scan), and early surgical intervention are of paramount importance to achieve a favourable outcome. Although the optimal therapeutic procedure for this entity is still a matter of discussion, enterolithotomy alone has been advocated as first-line approach for the majority of patients due to its lower morbidity, mortality, and reports on spontaneous fistula closure.^{2 3}

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Bär F**, Roblick U, Lehnert H. Gallstone ileus. *Clin Gastroenterol Hepatol* 2011;**9**:A22.
2. **Ayantunde AA**, Agrawal A. Gallstone ileus: diagnosis and management. *World J Surg* 2007;**31**:1292–7.
3. **Reisner RM**, Cohen JR. Gallstone ileus: a review of 1001 reported cases. *Am Surg* 1994;**60**:441–6.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Carrascosa MF, Riego-Martín MD, Cavedes JRS, Gutiérrez PG. Gallstone ileus. *BMJ Case Reports* 2012;10.1136/bcr.12.2011.5387, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow

Keep up to date with all published cases by signing up for an alert (all we need is your email address) <http://casereports.bmj.com/cgi/alerts/etoc>