Images in...

Wormy surprise!

Abraham Vincent Samuel Jesudoss, Meryem Kaya, Selina Lisa, Ashish Rohatgi

General Surgery Department, Whipps Cross University Hospital, London, UK

Correspondence to Mr Abraham Vincent Samuel Jesudoss, abepfdr@gmail.com

DESCRIPTION

A 44-year-old male presented with history of vomiting, abdominal pain and weight loss of 2 months duration. On examination, he was emaciated and had fullness in the epigastric region. Oesophagogastroduodenoscopy showed pyloric stenosis and biopsies showed mixed infiltrate of chronic inflammatory cells with no dysplasia and *Helicobacter pylori* organisms. CT scan showed thickening of the antro-pyloric region with dilatation of the stomach. In view of gastric outflow obstruction, he underwent partial gastrectomy and roux-en-y gastro-jejunostomy. During gastrojejunostomy, 7 metres tape worm was extracted from jejunal enterotomy (figure 1 and video 1). Histopathology showed granulation tissue and fibrosis within the pylorus and duodenal wall. No features of malignancy was evident. Postoperatively the patient was treated with praziquantel. Tapeworms are parasites that belong to Class cestoda and known as cestodes. There were about 71 cases of infestation with beef and pork tapeworms in UK and wales in 2005.¹ Since our patient does not consume pork, the cestode was likely to be beef tapeworm (*Taenia saginata*) and it is the most common tapeworm to infest humans. Transmission is via undercooked beef with cysticerci. Patients experience symptoms like nausea, vomiting, diarrhoea, abdominal discomfort, constipation and headache, dizziness or convulsions if neurocysticercosis present. Surgical complications like colonic perforation,² intestinal obstruction³ and appendicitis⁴ due to tapeworms have been reported. The diagnosis can be made by investigations like stool microscopy, detecting *Taenia* eggs in the perianal region and immunoblot assay to detect IgG anticysticercal antibodies in serum, cerebrospinal fluid and saliva.⁵ Anthelmintics like praziquantel and niclosamide are effective for treating intestinal tapeworm infestation.^{6,7} Albendazole is used for treating neurocysticercosis.

Video 1 Extraction of tapeworm via jejunal enterotomy.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. NHS Choices. Tapeworm infections. <http://www.nhs.uk/Conditions/Tapeworm-infections/Pages/Introduction.aspx>. (accessed 15 December 2011).
2. Demiriz M, Günhan O, Celasun B, *et al*. Colonic perforation caused by taeniasis. *Trop Geogr Med* 1995;**47**:180–2.
3. Bordon LM. Intestinal obstruction due to *Taenia saginata* infection: a case report. *J Trop Med Hyg* 1992;**95**:352–3.
4. Aydın O. Incidental parasitic infestations in surgically removed appendices: a retrospective analysis. *Diagn Pathol* 2007;**2**:16.

Figure 1 Tapeworms in kidney dish.

5. **Oliveira HB**, Machado GA, Mineo JR, *et al*. Taenia saginata metacestode antigenic fractions without affinity to concanavalin A are an important source of specific antigens for the diagnosis of human neurocysticercosis. *Clin Vaccine Immunol* 2010;**17**:638–44.
6. **Craig P**, Ito A. Intestinal cestodes. *Curr Opin Infect Dis* 2007;**20**:524–32.
7. **Garcia HH**, Del Brutto OH, Nash TE, *et al*. New concepts in the diagnosis and management of neurocysticercosis (Taenia solium). *Am J Trop Med Hyg* 2005;**72**:3–9.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Jesudoss AVS, Kaya M, Lisa S, Rohatgi A. Wormy surprise!. *BMJ Case Reports* 2012;10.1136/bcr.12.2011.5416, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow

Keep up to date with all published cases by signing up for an alert (all we need is your email address) <http://casereports.bmj.com/cgi/alerts/etoc>