Successful endovascular treatment of iatrogenic pseudoaneurysm with a relatively long neck

Masamichi Koganemaru,¹ Toshi Abe,¹ Naofumi Hayabuchi,¹ Kanako Yokoi²

¹Department of Radiology, Kurume University School of Medicine, Kurume, Japan

²Department of Internal Medicine, Kurume University School of Medicine, Kurume, Japan

Correspondence to

Dr Masamichi Koganemaru, mkoganemaru@med.kurume-u.ac.jp

DESCRIPTION

A 63-year-old man was admitted to our emergency medical centre due to suspected myocardial infarction; coronary angiography was performed via the left common femoral artery. Following cardiac intervention, colour doppler ultrasonography demonstrated left common femoral artery pseudoaneurysm. (CT) revealed a pseudoaneurysm sac (maximum size, 54 mm) with a relatively long neck (length, 27 mm; figure 1).

After failed ultrasound-guided compression, left external iliac arteriogram confirmed that the pseudoaneurysm sac communicates via a neck with the left common femoral artery (figure 2A). A 2 Fr-microcatheter was superselectively catheterised into the pseudoaneurysm neck (figure 2B), followed by embolisation by injecting a 1:1 mixture of n-butyl cyanoacrylate (NBCA) and lipiodol to enable fluoroscopic visualisation. Left external iliac arteriogram confirmed complete absence of the pseudoaneurysm immediately after embolisation (figure 2C). Follow-up CT imaging 12 days later revealed no traces of the pseudoaneurysm. Moreover, there were no complications associated with transcatheter embolisation.

Femoral artery pseudoaneurysms occur in 0.2–7.7% of patients undergoing angiographic procedures.¹ In general, coil embolisation is an effective treatment at sites both distal and proximal to the pseudoaneurysm.² However, in the present case, the technique may have caused lower limb

ischaemia. Treatment options for pseudoaneurysm include ultrasound-guided compression and direct percutaneous thrombin injection.³ NBCA is a liquid embolic material, and the adhesion time can be flexibly adjusted according to the rate of mixed lipiodol. A distinct advantage of NBCA in lipiodol is its dense radiopacity; as the exact site of occlusion can be observed during the embolisation procedure. NBCA undergoes rapid polymerisation and solidification and provides permanent embolisation; hence, migration of the embolic material can be prevented. The present technique demonstrated that superselective embolisation of the pseudoaneurysm neck was possible without the need for direct percutaneous embolisation.

Learning points

- ▶ Iatrogenic pseudoaneurysm is an undesirable complication of angiography.
- ▶ Endovascular treatment for pseudoaneurysm should be effective, particularly with a relatively long neck.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Figure 1 Contrast-enhanced CT in the arterial phase revealed a well-enhanced area (arrowhead) in the left inguinal region (A). Three-dimensional CT revealed a common femoral artery pseudoaneurysm (arrowhead) with a relatively long neck (arrow) (B).

To cite: Koganemaru M, Abe T, Hayabuchi N, *et al.* *BMJ Case Reports* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-007021

Figure 2 Left external angiogram revealed a pseudoaneurysm sac (arrowhead) and neck (arrow) (A). A microcatheter was superselectively catheterised into the pseudoaneurysm neck (thin arrow), followed by embolisation injecting a mixture of n-butyl cyanoacrylate and lipiodol (B). Left external iliac arteriogram immediately after embolisation confirmed complete absence of the pseudoaneurysm (C).

REFERENCES

- 1 Nakai M, Sato M, Sanda H, *et al*. Percutaneous fluoroscopically guided n-butyl cyanoacrylate (NBCA) injection for iatrogenic femoral arterial pseudoaneurysm under temporary balloon occlusion of arterial blood flow. *Jpn J Radiol* 2012;30:365–9.
- 2 Ikeda O, Nakasone Y, Tamura Y, *et al*. Endovascular management of visceral artery pseudoaneurysms: transcatheter coil embolization using the isolation technique. *Cardiovasc Interv Radiol* 2010;33:1128–34.
- 3 Chatterjee T, Do DD, Kaufmann U, *et al*. Ultrasound-guided compression repair for treatment of femoral artery pseudoaneurysm: acute and follow-up results. *Cathet Cardiovasc Diagn* 1996;38:335–40.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow