Ventricular ectopy after exercise and occult ischaemia in a patient with multiple risk factors for ischaemic heart disease and defective anginal warning system

Gian Piero Carboni,¹ Pietro Sedati²

¹Department of Nuclear Cardiology, Università Campus Bio-Medico, Rome, Italy
²Department of Radiology, Università Campus Bio-Medico, Roma, Italy

Correspondence to
 Professor Gian Piero Carboni,
 g.carboni@unicampus.it

DESCRIPTION

In May 2012, a 73-year-old man presented with effort dyspnoea and chest discomfort, negative exercise test for ST depression and ventricular ectopy after exercise. He had a history of Sjögren's syndrome, diabetes, dyslipidemia, hypertension, chronic obstructive pulmonary disease and was under methotrexate treatment for rheumatoid arthritis. Such diseases imply a severe dysfunctional burden, tend to develop accelerated atherosclerosis, endothelial dysfunction and ischaemic heart disease (IHD).¹⁻² A functional assessment was thus provided with an exercise/rest technetium-99 m tetrofosmin-gated single-photon emission cardiac

tomography (G-SPECT). This test did not show perfusion defects; left ventricular function was normal (figure 1). At peak exercise, there was lack of ST-depression, but episodes of ventricular triplets were detected after exercise (figure 2). A cardiac tomography (CT) scan detected a severe stenosis in the middle portion of the left anterior descending (LAD) coronary artery (figure 3). An invasive coronary angiography revealed 90% stenosis of the LAD; therefore, a stent was implanted. Ventricular ectopy after exercise is associated with reactivation of parasympathetic activity and an increased risk of death.³ In the presence of a subocclusive LAD stenosis, the patient's exercise SPECT results should be thus

Figure 1 Exercise-gated /rest-gated single-photon emission cardiac tomography shows lack of significant perfusion defects and a normal left ventricular function.

To cite: Carboni GP, Sedati P. *BMJ Case Reports* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-007536

Figure 2 A run of ventricular triplets and an isolated ventricular ectopy during the exercise test recovery period. Heart rate (HR); beats per minute (bpm) and systolic blood pressure (SBP).

considered as false negative. Occult exercise-induced ischaemia and a reperfusion mechanism for postexercise ventricular ectopy cannot be ruled out. Reperfusion re-establishes slow conduction through depressed regions, permitting re-entrant pathways to form again, resulting in the re-emergence of ventricular arrhythmias.⁴ Cardiac CT may provide early-stage screening for occult IHD in patients with multiple risk factors for IHD and defective anginal warning system.

Figure 3 Volume-rendered cardiac CT images indicate a severe stenosis in the middle portion of the anterior descending (LAD) artery. Aorta(Ao), pulmonary trunk (PT) and left circumflex artery (Cx).

Learning points

- ▶ Rheumatoid arthritis and other rheumatic diseases significantly increase the risk of atherosclerosis.
- ▶ Diabetes, rheumatoid arthritis and Sjögren's syndrome may lead to uncommon presentations of ischaemic heart disease.
- ▶ If pretest probability is high, negative test results should not stop investigations.
- ▶ Ventricular arrhythmias may be a sign of cardiac hypoperfusion even in a negative test.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Mellana WM, Aronow WS, Palaniswamy C, *et al*. Rheumatoid arthritis: cardiovascular manifestations, pathogenesis, and therapy. *Curr Pharm Des* 2012;18:1450–6.
- 2 Inoue H, Kinoshita K, Sugiyama M, *et al*. Sudden death from ischaemic heart disease in a female patient with Sjögren syndrome: a case report. *Med Sci Law* 2008;48:261–5.
- 3 Frolkis JP, Pothier CE, Blackstone EH, *et al*. Frequent ventricular ectopy after exercise as a predictor of death. *N Engl J Med* 2003;348:781–90.
- 4 Murdock DK, Loeb JM, Euler DE, *et al*. Electrophysiology of coronary reperfusion. A mechanism for reperfusion arrhythmias. *Circulation* 1980;61:175–82.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow