Missing clues for a radiologist

Tushar Ramesh Sahasrabudhe

Department of Pulmonary
Medicine, Padmashree Dr. D.Y.
Patil Medical College, Pune,
Maharashtra, India

Correspondence to
Professor Tushar Ramesh
Sahasrabudhe,
drtrs@sify.com

DESCRIPTION

A 48-year-old man presented with predominant nocturnal cough with minimal sputum, since 2 years. He had frequent retrosternal pain with occasional lower dysphagia since 6 months. He had no weight loss, fever or wheeze. He was a hypertensive for 9 years. He had left leg fracture at the age of 11 and developed permanent limb shortening in spite of multiple surgeries. He had mild cognitive defect since 1 year.

His chest x-ray (figure 1) showed a large retrocardiac globular shadow without a fluid level, unclear borders of left hemi diaphragm and multiple pulmonary nodules. The radiological diagnosis given by the radiologist was a posterior mediastinal mass with multiple lung metastases, most likely a lower oesophageal malignancy.

The reported large mediastinal mass and metastases were actually the tumours on his chest wall (figures 2 and 3). The patient is suffering from Neurofibromatosis type I (NF1), also known as Von Recklinghausen disease.¹ The chest radiogram was reported by a radiologist, who neither saw the patient personally nor enquired into the patient's history. Note the multiple café au lait spots on the patient's skin, another characteristic of this disease.² Cognitive defects, pseudoarthrosis of tibia and long bone dysplasia are also known to occur in NF1.³ Pheochromocytoma was not present in him but is known in NF1.

The patient underwent oesophago-gastroscopy which confirmed the presence of GERD (Gastro-oesophageal reflux disease). Oesophageal

Figure 1 Chest x-ray posteroanterior view.

Figure 2 Large chest wall tumour on the lower end of sternum.

malignancy or hiatus hernia was ruled out. The cough responded to antireflux treatment within 1 month.

Figure 3 Posterior chest wall tumours.

To cite: Sahasrabudhe TR.
BMJ Case Reports Published
online: 2 January 2013
doi:10.1136/bcr-2012-
007715

Learning points

- ▶ Soft tissue lesions on the anterior or posterior chest wall can be easily mistaken as intrapulmonary lesions on a chest radiogram posteroanterior (PA) view.
- ▶ A standard chest radiogram PA view should always cover the lateral chest wall. Had this been done, the skin nodules would have been easily picked up.
- ▶ A clinician should always clinically correlate an x-ray and its report, before making a diagnosis.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Rubin R, Strayer DS. *Rubin's pathology: clinicopathologic foundation of medicine*. 5th edn. Baltimore: Wolters Kluwer Health, Lippincott Williams & Wilkins, 2008:201–3.
- 2 Torpy JM, Burke AE, Glass RM. JAMA patient page: neurofibromatosis. *JAMA* 2008;300:352.
- 3 Riccardi VM. Von Recklinghausen neurofibromatosis. *N Engl J Med* 1981;305:1617–26.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow