Radiographical appearance of osteitis fibrosa cystica in primary hyperparathyroidism before and after parathyroidectomy

Mark Anthony Santiago Sandoval

Section of Endocrinology,
Diabetes and Metabolism,
Department of Medicine,
College of Medicine and
Philippine General Hospital,
University of the Philippines
Manila, Manila, Philippines

Correspondence to

Dr Mark Anthony Santiago Sandoval,
markanthony_sandoval@
yahoo.com

DESCRIPTION

The case of a patient with primary hyperparathyroidism with classic and severe skeletal involvement has been previously reported.¹ Briefly, it was the case of a 63-year-old woman who presented with multiple repeated fractures, the latest of which were of both femoral shafts. Primary hyperparathyroidism was diagnosed based on elevated calcium with unsuppressed parathyroid hormone (PTH) levels. The excess PTH secretion was localised to a right inferior parathyroid adenoma. She eventually underwent parathyroidectomy with documented decline in calcium and PTH levels.

A preoperative skeletal survey radiograph was performed on November 2008, 3 months before parathyroidectomy. Among the findings was a lytic lesion of the fourth metacarpal of the right hand, compatible with osteitis fibrosa cystica. Repeat x-ray performed on April 2012 or 38 months after parathyroidectomy showed complete remineralisation of the previously noted lytic lesion. The area previously occupied by the osteitis fibrosa cystica now appears hyperdense indicative of bone formation (figure 1).

Her fractures have also healed and there has been no recurrence of any fracture since the surgery. She has also been ambulatory since then.

Areas of the bone affected by osteitis fibrosa cystica are known to resolve, with radiographs showing hyperdensities within 3 months of successful parathyroidectomy.² These radiographical changes are also accompanied by histological changes in bone, showing an osteoblastic or healing response.³

Learning points

- ▶ Osteitis fibrosa cystica pertains to lytic lesions of the bone that appear as cysts. These are due to excessive bone resorption seen in primary hyperparathyroidism.
- ▶ Osteitis fibrosa cystica resolves after parathyroidectomy, with radiographs showing complete remineralisation and bone formation.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

November 2008

3 months before intervention

April 2012

38 months after intervention

Figure 1 On a radiograph prior to parathyroidectomy (the intervention), a lytic lesion was seen on the fourth metacarpal of the right hand (yellow arrow), compatible with osteitis fibrosa cystica. On follow-up performed 38 months after parathyroidectomy, there was now a hyperdensity occupying the same area indicating complete remineralisation of the lytic lesion (blue arrow).

REFERENCES

- 1 Sandoval MAS, Paz-Pacheco E. Primary hyperparathyroidism with classic and severe skeletal involvement. *BMJ Case Rep* 2010. doi:10.1136/bcr.04.2010.2929
- 2 Agarwal G, Mishra SK, Kar DK, *et al*. Recovery pattern of patients with osteitis fibrosa cystic in primary hyperparathyroidism after successful parathyroidectomy. *Surgery* 2002;132:1075–83; discussion 1083–5.
- 3 Chavda DV, Frock JT, Zielinski CM, *et al*. Reversal of histology of bone after parathyroidectomy in patients with hyperparathyroidism. *J South Orthop Assoc* 1998;71: 65–71.

To cite: Sandoval MAS.
BMJ Case Reports Published
online: [please include Day
Month Year] doi:10.1136/
bcr-2012-008086

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow