CASE REPORT

Giant gastric folds in a patient with hypoalbuminaemia

Jurgen Gerada,¹ Elaine Borg,² James DeGaetano,² James Pocock¹¹Division of Gastroenterology, Mater Dei Hospital, Msida, Malta²Department of Histopathology, Mater Dei Hospital, Msida, Malta**Correspondence to**
Dr Jurgen Gerada,
jurgen.gerada@gmail.com**DESCRIPTION**

A 70-year-old man presented with a 3-month history of persistent epigastric pain, 8 kg weight loss and lower limb swelling. Examination revealed a normal abdomen and bilateral lower limb oedema. Blood results were unremarkable except for hypoalbuminaemia (23 g/l) and peripheral eosinophilia ($1.06 \times 10^9/l$). Urinalysis and CT of the abdomen were negative. An oesophagogastroduodenoscopy (OGD) revealed large gastric folds involving the fundus and the body of the stomach (figure 1), and a small antral polyp. Histology showed marked reactive mucosal changes, pronounced foveolar hyperplasia and cystic dilatation (figure 2), compatible with a diagnosis of Ménétrier's disease. *Helicobacter pylori* was absent and the antral polyp was adenomatous. He was given omeprazole and prednisolone (20 mg daily), which was tailed down after 2 months. Subsequently, his symptoms improved significantly. He regained 10 kg in weight and his albumin levels normalised. OGD done post therapy and 1 year later still showed prominent, but less pronounced, gastric folds, and histological evidence of hyperplastic gastropathy.

The current treatment options to manage Ménétrier's disease include testing and treating cytomegalovirus and *H pylori* infection, proton pump inhibitor or histamine-2 receptor blocker, octreotide, monoclonal antibodies to epidermal growth factor receptor (EGFR) and gastrectomy.¹ Corticosteroid use in this setting has been reported very rarely, successfully in a 69-year-old male patient,² but not so in a 4-year-old boy.³ To our knowledge, our case is the

Figure 2 Hyperplastic gastropathy showing pronounced foveolar hyperplasia.

second case to have responded to steroids. Hence, one might consider this treatment before other expensive treatments like anti-EGFR therapy.

Learning points

- ▶ Ménétrier's disease is a rare condition and a high level of suspicion should be maintained when a patient presents with upper gastrointestinal symptoms, hypoalbuminaemia and giant gastric folds at endoscopy.
- ▶ Ménétrier's disease affects men three times as much as women and is known to predispose to gastric carcinoma.
- ▶ Corticosteroid use might be an alternative to treat this condition.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Soares JB, Bastos P, Goncalves R. Ménétrier disease with antrum polyposis and gastritis cystica profunda. *Endoscopy* 2012;44(Suppl 2 UCTN):E56–7.
- 2 Winney RJ, Gilmour HM, Matthews JD. Prednisolone in giant hypertrophic gastritis (Menetrier's disease). *Am J Dig Dis* 1976;21:337–9.
- 3 Ling LJ, Hershenson MB, Young S, et al. Home parenteral nutrition in a child with Menetrier disease. *Eur J Pediatr* 1986;144:505–7.

Figure 1 Giant gastric folds of the body of the stomach.

To cite: Gerada J, Borg E, DeGaetano J, et al. *BMJ Case Reports* Published Online: [please include Day Month Year] doi:10.1136/bcr-2012-008284

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow