String test

Yoshiro Hadano

Department of General Internal Medicine and Infectious Diseases, Rakuwakai Otowa Hospital, Kyoto, Japan

Correspondence to

Dr Yoshiro Hadano,
hatayoshiyoshi@gmail.com

DESCRIPTION

An 82-year-old Japanese woman with duodenum papilla cancer was admitted with a 3-day history of fever, chills and nausea. A contrast-enhanced CT scan showed ring-enhanced and multiloculated low-density area in her liver and suspected pyogenic liver abscess (figure 1).

Drainage of the liver abscess was performed and Gram's staining of the abscess revealed Gram-negative bacilli with capsule formation. Blood cultures and abscess cultures grew *Klebsiella pneumoniae*. Growing colonies from the liver abscess on blood agar were shiny, cream-coloured, and the string test was positive (>5 mm string; figure 2). The patient was treated with cefepime for 2 weeks and switched to oral amoxicillin/clavulanate for an additional 4 weeks. No evidence of relapse of the infection was noted at the 6-month follow-up.

Community-acquired *K pneumoniae* liver abscess sometimes causes metastatic region such as ocular or central nervous system, and hypermucoviscous phenotype of *K pneumoniae* is associated with hypervirulence. The presence of capsular serotype K1, and to a lesser extent K2, appears to play a

Figure 2 Demonstrating the string test.

role in the virulence of this organism,^{1 2} and it is easy and very useful to identify the hypervirulence type of *K pneumoniae* by a string test in the community hospital's laboratory.¹

Learning points

- ▶ Hypermucoviscous phenotype of *Klebsiella pneumoniae* is associated with hypervirulence.
- ▶ It is easily identified by a string test in the community hospital's laboratory without Caspar serotyping system.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Fang CT, Chuang YP, Shun CT, *et al.* A novel virulence gene in *Klebsiella pneumoniae* strains causing primary liver abscess and septic metastatic complications. *J Exp Med* 2004;199:697–705.
- 2 Fang CT, Lai SY, Yi WC, *et al.* *Klebsiella pneumoniae* central nervous system complications from pyogenic liver abscess. *Clin Infect Dis* 2007;45:284–93.

Figure 1 Contrast-enhanced CT scan showing findings suggestive of liver abscess.

To cite: Hadano Y. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-008328

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow