Basic and advanced imaging of a case of Balò's concentric sclerosis

Paolo Ripellino,^{1,2} Alessandro Stecco,³ Maurizio A Leone,² Roberto Cantello²

¹Department of Neurology, University of Turin, Torino, Italy
²Department of Neurology, University of Eastern Piedmont, AOU Maggiore della Carità, Novara, Italy
³Department of Radiology, University of Eastern Piedmont, AOU Maggiore della Carità, Novara, Italy

DESCRIPTION

In January 2012 a 51-year-old woman was admitted to our department for a slow increasing right hemiparesis with right brisk deep tendon reflexes. She suffered from Sjogren syndrome, fibromyalgia and hepatitis B virus (healthy carrier).

Conventional MRI scans (figure 1A,B) were typical for Balò's sclerosis.¹

The patient rejected a brain biopsy. In order to better define the lesion, we therefore performed advanced radiological techniques such as diffusion, spectroscopy and perfusion.

In the diffusion sequences (figure 2A) a concentric ring of unrestricted diffusion appeared clearly distinguishable.²

The spectroscopic image (figure 2B) of the lesion documented increased choline peak and decreased N-acetyl aspartate peak, but normal peaks in the normal appearing white matter near the lesion.¹

The perfusion analysis (figure 2C), based on four different regions of interest (figure 2D), suggested a decreasing gradient of perfusion from the centre to the periphery of the lesion, supporting the hypothesis that the centre of the ring corresponds anatomically to a deep venous vessel.³ Authors are not aware of previous report of perfusion studies in Balò's sclerosis.

Cerebrospinal fluid analysis detected oligoclonal bands, without anti-aquaporin-4 antibodies.

Figure 1 At diagnosis. (A) Axial fluid attenuated inversion recovery sequence and (B) sagittal T2-sequence evidence the typical concentric ring lesion in the white matter.

Figure 2 At diagnosis. (A) Diffusion shows a restricted peak depicting the classical ring of demyelination and remyelination. (B) Spectroscopy shows a high ratio Cho/N-Acetylaspartate, double pick of lactate-lipids. (C) Perfusion MRI analysis, based on four regions of interest depicted in D (A, pink, B, white, C, blue, D, red) suggests that the central layers of the ring are more perfused compared to the peripheral ones.

To cite: Ripellino P, Stecco A, Leone MA, et al. *BMJ Case Reports* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-008413

Figure 3 At 8 months follow-up (after treatment). (A) Axial fluid attenuated inversion recovery sequence and (B) sagittal T2-sequence evidence the reduction in the lesion volume compared to figure 1.

High-dose intravenous methylprednisolone was started (1 g/day for 10 days) and followed by oral prednisone (1 mg/kg/day for 2 months, then slowly tapered in 2 months) with concomitant lamivudine as antiviral prophylaxis. The right paresis fully recovered after 2 weeks and has not relapsed after 11 months.

The 8 months follow-up MRI showed a reduction in the lesion volume and absence of new demyelinating lesions (figure 3A,B).

Learning points

- ▶ Baló's concentric sclerosis is a rare demyelinating disease presenting with a concentric ring in the white matter. The clinical and radiological features can mimic other diseases such as primary central nervous system lymphoma, low-grade glioma or stroke.
- ▶ A brain biopsy should be obtained whenever possible. However, nowadays advanced neuroimaging studies (spectroscopy, MRI diffusion and perfusion) seem to be a reliable tool for the diagnosis.
- ▶ Baló's sclerosis could respond very well to high-dose steroids alone or in combination with other immunosuppressive treatments (eg, plasma exchange).

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Karaarslan E, Altintas A, Senol U, *et al*. Baló's concentric sclerosis: clinical and radiologic features of five cases. *AJNR Am J Neuroradiol* 2001;22:1362–7.
- 2 Kavanagh EC, Heran MK, Fenton DM, *et al*. Diffusion-weighted imaging findings in Baló concentric sclerosis. *Br J Radiol* 2006;79:e28–31.
- 3 Khonsari RH, Calvez V. The origins of concentric demyelination: self-organization in the human brain. *PLoS One* 2007;2:e150.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow