Enthesitis in association with inflammatory bowel disease

Iain Edward Ewing,¹ Eleni Josephides,² Daniel Galloway,² Pranab Gyawali²

¹Department of Gastroenterology, North Middlesex Hospital, London, UK

²Department of Gastroenterology, Queen's Hospital, BHR Hospitals NHS Trust, London, UK

Correspondence to

Dr Iain Edward Ewing,
iainewing@doctors.org.uk

DESCRIPTION

A 27-year-old Sudanese woman presented to the emergency department with a 1-week history of lumbar back pain radiating to the buttocks. There was no history of trauma, fever or weight loss. Medical history included ulcerative colitis, maintained in remission with oral mesalazine. On examination, there was midline tenderness over the upper lumbar spine. Neurological examination was normal.

MRI of the lumbar spine (figure 1) demonstrated oedema in the subcutaneous tissue adjacent to the first lumbar (L1) vertebra and bone marrow oedema of the corresponding spinous process. These changes were reported as being consistent with contusion secondary to trauma. This was incompatible with the clinical history.

While undergoing further evaluation, the patient developed acute bloody diarrhoea. Stool microscopy, culture and *Clostridium difficile* toxin were negative. Flexible sigmoidoscopy showed severely active ulcerative colitis with spontaneous bleeding and ulceration.

In the context of active ulcerative colitis, the MRI was reviewed again. The appearances are consistent with enthesitis of the supraspinous ligament, maximal at the point of insertion at the L1 spinous process, with a differential diagnosis not initially considered.

Treatment was initiated with intravenous hydrocortisone. Symptoms of enthesitis and ulcerative colitis both responded rapidly. Azathioprine was given alongside a reducing course of corticosteroids to maintain long-term remission.

Enthesitis is a component of seronegative spondyloarthropathy associated with inflammatory bowel disease (IBD).¹ The prevalence of spondyloarthropathy was reported to be 22% in a population study of IBD patients.² Disease activity of the two entities is not always related.³

Learning points

- ▶ If a clinically unexplained finding is reported on the diagnostic imaging, revisit the history and explore the differential diagnosis.
- ▶ Remember to consider known disease associations.
- ▶ Seronegative arthropathy is commonly associated with inflammatory bowel disease (IBD) and should be considered in IBD patients presenting with rheumatological symptoms.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 McGonagle D, Stockwin L, Isaacs J, *et al.* An enthesitis based model for the pathogenesis of spondyloarthropathy: additive effects of microbial adjuvant and biomechanical factors at disease sites. *J Rheumatol* 2001;28:2155–9.
- 2 Palm O, Moum B, Ongre A, *et al.* Prevalence of ankylosing spondylitis and other spondyloarthropathies among patients with inflammatory bowel disease: a population study (the IBSEN study). *J Rheumatol* 2002;29:511–15.
- 3 Brakenhoff LK, van der Heijde DM, Hommes DW. IBD and arthropathies: a practical approach to its diagnosis and management. *Gut*;60:1426–35.

Figure 1 T2-weighted MRI of the lumbar spine demonstrating oedema in the subcutaneous tissue adjacent to the first lumbar (L1) vertebra and bone marrow oedema of the L1 spinous process.

To cite: Ewing IE, Josephides E, Galloway D, *et al.* *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-008418

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow