Phthiriasis palpebrarum

Georgios D Panos,¹ Ioannis K Petropoulos,² Doukas Dardabounis,³ Zisis Gatzoufas¹

¹Department of Ophthalmology, University Hospitals of Geneva, Geneva, Switzerland

²Department of Ophthalmology, Ophthalmological Center of Rive, Geneva, Switzerland

³Department of Ophthalmology, University General Hospital of Alexandroupolis, Alexandroupolis, Eastern Macedonia & Thrace, Greece

Correspondence to

Dr Georgios D Panos, gdpanos@gmail.com

DESCRIPTION

A 61-year-old woman was admitted to the hospital because of itching, burning sensation and moderate pain of 3 months duration in her right eye. Her ophthalmologist diagnosed common anterior blepharitis and prescribed a combination of antibiotic and corticosteroid eye drops. Best corrected visual acuity was 20/20 in both eyes, intraocular pressure was 15 mm Hg right eye and 14 mm Hg left eye. Slit lamp examination revealed nits anchored to the eyelashes (figure 1). Moving insects were observed hanging from the eyelashes (red arrow, figure 1). We mechanically removed some of the eyelashes, including lice and nits, with the use of fine forceps and advised the patient to apply yellow oxide of mercury ointment 1% three times daily over the lid margins. Ten days later, the eyelashes were clear of all lice and nits. We continued the same treatment once daily for about 3 weeks. The patient was followed-up once every 15 days for 2 months. There was complete cure without any recurrence.

Phthiriasis palpebrarum is an uncommon condition in developed countries and appears in cases of bad hygiene. Sometimes this disease can be misdiagnosed as common blepharitis. There are a number of treatment options for phthiriasis palpebrarum including

cryotherapy, argon laser photocoagulation, fluorescein eye drops 20%, physostigmine 0.25%, lindane 1%, petroleum gel, yellow mercuric oxide ointment 1%, malathion drops 1% or malathion shampoo 1%, oral ivermectin and pilocarpine gel 4%.¹ However, the most popular and simplest treatment is direct removal of the parasites with forceps.^{1–3} Family members, sexual and close contacts should all be examined and treated appropriately, if necessary.

Learning points

- ▶ Phthiriasis palpebrarum is an uncommon condition in developed countries and appears in cases of bad hygiene.
- ▶ Sometimes this disease can be misdiagnosed as common blepharitis.
- ▶ There are a number of treatment options for phthiriasis palpebrarum, however the most popular and simplest is the direct removal of parasites with fine forceps.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Jiang J, Shen T, Hong CY. A peculiar case of eye pruritus: phthiriasis palpebrarum initially misdiagnosed as common blepharitis. *Int J Ophthalmol* 2011;4:676–7.
- 2 Perlman HH, Fraga S, Medina M. Phthiriasis palpebrarum (phthiriasis ciliaris). *J Pediatr* 1956;49:88–90.
- 3 Ronchese F. Treatment of pediculosis ciliarum in an infant. *N Engl J Med* 1953;249:897–8.

Figure 1 Slit lamp examination: Nits anchored to the eyelashes. A moving insect was observed hanging from an eyelash (red arrow).

To cite: Panos GD, Petropoulos IK, Dardabounis D, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-009272

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>. BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow