Large spontaneous pneumothorax with relatively minor symptoms

Tamsin E M Morrison

Brighton and Sussex University
Hospitals NHS Trust, East
Sussex, Brighton, UK

Correspondence to
Dr Tamsin Morrison, tamsin.
morrison@doctors.org.uk

DESCRIPTION

A 40-year-old gentleman presented to emergency department (ED) with sudden onset of right-sided chest pain, largely subsided on admission. He was an ex-smoker with a history of basal cell carcinoma and patella fracture. Trachea was central on examination, with absence of right-sided breath sounds. Observations were normal.

Chest radiograph revealed a large right-sided primary spontaneous pneumothorax (PSP) (figure 1), which was aspirated unsuccessfully. A chest drain was inserted with uneventful recovery (figure 2).

Global occurrence is greater in men, at 0.018–0.028% versus 0.0012–0.006%, with 54% risk of another pneumothorax within 4 years.¹ Smoking poses a further 12% risk.¹ Alveoli of taller subjects are also prone to greater distension pressure and development of subpleural blebs as negative pleural pressure gradient increases towards the lung apex.¹

If present, signs include reduced lung expansion, absent breath sounds and hyper-resonance on the affected side. Although diagnosis and management should be guided clinically, an inspiratory chest x-ray is recommended.¹ CT scanning, 'gold standard' for size estimation and detection of small pneumothoraces, is advised for uncertain cases.¹

Treatment depends on:

1. Previous lung pathology.
2. Breathlessness.
3. Pneumothorax size.
 - A. Insert chest drain if bilateral or haemodynamically unstable spontaneous pneumothoraces.

Figure 1 Erect anteroposterior chest radiograph showing large right-sided pneumothorax with almost complete right lung collapse with loss of right lung markings. There is no tracheal deviation. Pneumothorax defined as 'large' by the presence of greater than 2 cm rim between lung margin and chest wall.¹

Figure 2 Erect posteroanterior chest radiograph showing lung fields after chest drain removal. There is no pneumothorax present; however, there is some persistent atelectasis in the right lung. The left lung is clear.

- B. If age >50, clinical/radiological lung pathology or significant smoking history, treat as secondary pneumothorax.¹
- C. If PSP >2 cm or shortness of breath, aspirate less than 1.5 l. If aspiration fails, insert chest drain.
- D. If <2 cm with no shortness of breath, or successful aspiration, consider discharge with 2–4 week follow-up.¹

Learning points¹

- ▶ Spontaneous pneumothorax should always be considered in a young, tall, male patient who presents with sudden onset chest pain. Delayed presentation comes with the risk of re-expansion pulmonary oedema.
- ▶ Size of pneumothorax does not necessarily correlate with severity of symptoms—symptoms may even be absent in primary spontaneous pneumothorax and thus a high level of clinical diagnostic suspicion is required. If there are signs of cardio-respiratory compromise, do not delay treatment to perform imaging. Treat as a tension pneumothorax.
- ▶ Smoking increases the risk of primary spontaneous pneumothorax by a further 12% compared with 0.1% of non-smokers. There is no evidence to suggest development of pneumothorax is associated with physical exertion. However, in view of increased risk of recurrence of pneumothorax in an affected individual, controllable risk factors such as smoking and progression to chronic obstructive pulmonary disease should be addressed.

To cite: Morrison TEM. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-009531

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCE

- 1 MacDuff A, Arnold A, Harvey J, on behalf of the BTS Pleural Disease Guidelines Group. Management of spontaneous pneumothorax: British Thoracic Society pleural disease guideline. BTS Guidelines. *Thorax* 2010;65(Suppl 2):ii18–31.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow