Pyopneumopericardium and empyema thoracis from perforated oesophageal cancer

Nitinan Chimparlee,¹ Monravee Tumkosit,² Kittichai Luengtaviboon,³ Pairaj Chattranukulchai⁴

¹Department of Biochemistry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

²Department of Radiology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

³Division of Cardiothoracic Surgery, Department of Surgery, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

⁴Division of Cardiology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Cardiac Center, King Chulalongkorn Memorial Hospital, Bangkok, Thailand

Correspondence to

Dr Pairaj Chattranukulchai, pairaj.md@gmail.com

DESCRIPTION

A 79-year-old Thai man with a history of untreated mid-oesophageal cancer presented with progressive dyspnoea and productive cough for 3 weeks. On examination, he had low-grade fever, tachypnoea, regular pulse rate at 110/min, blood pressure 120/78 mm Hg without definite pulsus paradoxus. Pulmonary examination showed mild deviation of trachea to the right, dullness on percussion and decreased breath sound at the left lung. He also had mildly distended neck veins, decreased cardiac sound and decreased apical impulse. ECG revealed sinus tachycardia with low QRS voltage without significant ST-T change. A large amount of generalised high-echoic pericardial fluid and evidences of early cardiac tamponade were demonstrated by transthoracic echocardiography.

Contrast-enhanced CT of the chest showed known circumferential oesophageal mass at the middle one-third of the oesophagus (black asterisk, figure 1A). A large amount of circumferential pericardial fluid (white asterisks, figure 1A–C) with small air bubbles (bold arrows, figure 1A–C) were detected. In addition, large loculated fluid collections in left hemithorax with air-fluid level and pleural enhancement (open arrows, figure 1A,C) were observed. Subsequent pericardiocentesis along with left intercostal drainage were performed, which revealed similar findings of large volume of frank, yellowish, foul-smelling pus (figure 2), consistent with pyopneumopericardium and empyema thoracis. On the basis of these findings, the patient underwent surgery with the presumptive diagnosis of the perforation of necrotic oesophagus leading to direct spread of infection to adjacent organs. Intraoperative finding confirmed the ruptured oesophagus at mid-thoracic level as a consequence

Figure 2 Frank, yellowish, foul-smelling pus from pericardiocentesis which appeared similar to empyema thoracis from intercostal drainage. Microbiological culture was positive for mixed organisms including *Pseudomonas aeruginosa*, *Streptococcus group D* and *Streptococcus group G*.

of local tumour invasion. He successfully underwent combined drainage of pus in pericardial space with pericardiectomy, decortication of left lung and resection of the thoracic oesophagus from just below aortic arch to oesophagogastric junction. Microbiological cultures of pus were positive for mixed organisms including *Pseudomonas aeruginosa*, *Streptococcus group D* and *Streptococcus group G*. Postoperative period was uneventful and he is currently in the restoration process of enteral alimentation before being discharged.

Perforation of oesophageal cancer is a serious circumstance that most often results from either diagnostic or therapeutic instrumentation, that is, stent placement.¹ Locally advanced cancer may lead to

Figure 1 Contrast-enhanced CT in axial (A), coronal (B) and sagittal views (C) show malignant oesophageal mass encasing the nasogastric tube at mid-oesophagus (black asterisk, A). A large amount of circumferential pericardial effusion (white asterisks, A–C) with small air bubbles (bold arrows, A–C) indicated pyopneumopericardium. Large loculated fluid collections in left hemithorax with air-fluid level and pleural enhancement of its thin wall (open arrows, A and C) represented empyema thoracis.

To cite: Chimparlee N, Tumkosit M, Luengtaviboon K, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-010053

spontaneous perforation of oesophagus which has high mortality rate.^{2 3} This allows progressive migration of the normal bacterial flora from the oesophagus to adjacent mediastinal organs include of the pericardial sac and pleural cavity leading to uncontrolled infection.⁴ Pyopneumopericardium and empyema thoracis are very rare, but recognised, complications which dictate prompt surgical therapy.

Learning points

- ▶ Perforation of oesophageal cancer is a serious circumstance that most often results from either diagnostic or therapeutic instrumentation.
- ▶ Advanced oesophageal cancer may lead to spontaneous perforation from local tumour invasion.
- ▶ Pyopneumopericardium and empyema thoracis are very rare complications as a result of progressive migration of the normal bacterial flora from the oesophagus to adjacent mediastinal organs.

Contributors NCand PC wrote the manuscript. MT, KL reviewed and revised the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Huber-Lang M, Henne-Bruns D, Schmitz B. Esophageal perforation: principles of diagnosis and surgical management. *Surg Today* 2006;36:332–40.
- 2 Reeder LB, DeFilippi VJ, Ferguson MK. Current results of therapy for oesophageal perforation. *Am J Surg* 1995;169:615–17.
- 3 Bhatia P, Fortin D, Inculet RI, *et al*. Current concepts in the management of esophageal perforations: a twenty-seven year Canadian experience. *Ann Thorac Surg* 2011;92:209–15.
- 4 Piatkowski R, Kochanowski J, Karpiński G, *et al*. Purulent pericarditis in patient with esophageal cancer presenting with cardiac tamponade. *J Emerg Med* 2011;40:671–3.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow