Ventriculopleural shunt CSF pseudocyst presenting as an anterior mediastinal mass

Peter Christian Thurlow, Chris Somerville, William Joseph Rusnak, Timothy J Mickus

Department of Radiology,
Allegheny General Hospital,
Pittsburgh, Pennsylvania, USA

Correspondence to
Dr Peter Christian Thurlow,
pthurlow@wpahs.org

DESCRIPTION

An asymptomatic middle-aged man presented for routine preoperative chest radiographs prior to the resection of a Masson's tumour of the frontal sinus. The chest radiographs revealed a large anterior mediastinal mass (figure 1). A CT scan of the chest was subsequently performed for further evaluation, demonstrating a cystic mass abutting the right anterior mediastinum with homogenous fluid attenuation (figure 2). A catheter could be seen extending from the soft tissues of the neck and coiling within the mass. Correlation with the head CT led to a diagnosis of pleural cerebrospinal fluid (CSF) pseudocyst associated with a ventriculopleural shunt. On further discussion with the patient, it was revealed that he had undergone ventriculopleural shunt placement for hydrocephalus at an outside institution following an episode of traumatic subarachnoid haemorrhage 20 years earlier. No further intervention was required, as the patient remained asymptomatic.

A CSF pseudocyst is a well-recognised complication of CSF diversion catheters that is most commonly associated with ventriculoperitoneal shunts.¹ The ventriculopleural shunt is an alternative method of CSF diversion used when ventriculoperitoneal or ventriculoatrial shunts are not suitable due to adhesions, a history of peritonitis, ascites, peritoneal dialysis or failure of a prior ventriculoperitoneal shunt.² Although uncommon, the ventriculopleural shunt is associated with a number of complications in the chest, including pleural effusion, empyema, tension hydrothorax, pneumothorax and rarely, as in this case, a CSF pseudocyst.² An understanding of this alternative method of CSF diversion and awareness of the potential complications is essential when radiographic abnormalities of the chest are encountered in a patient with a ventriculopleural shunt.

Figure 1 Posteroanterior (A) and lateral (B) radiographs of the chest demonstrate a rounded anterior mediastinal mass (asterisk). Differential considerations at the time of interpretation included lymphoma, thymoma or germ cell tumour.

CrossMark

To cite: Thurlow PC,
Somerville C, Rusnak WJ,
et al. *BMJ Case Rep*
Published online: [please
include Day Month Year]
doi:10.1136/bcr-2013-
201303

Figure 2 Contrast-enhanced CT of the chest demonstrating a ventriculopleural shunt catheter extending into a pleural cerebrospinal fluid (CSF) pseudocyst. (A) Axial images demonstrate a loculated pleural fluid collection (asterisk) measuring 0 hounsfield units, representing simple fluid density. (B) Coronal reconstructions showing a catheter (arrow) extending from the right neck into the pleural fluid collection. (C and D) Sagittal and three dimensional reconstructions with right anterior ribs digitally subtracted showing the catheter (arrowheads) coiling within the pleural fluid collection. A CT scan of the head (not shown) revealed a ventricular shunt in the right lateral ventricle with catheter tubing extending down the right side of the head. This constellation of findings led to the diagnosis of a pleural CSF pseudocyst as a complication of a ventriculopleural shunt.

Learning points

- The ventriculopleural shunt is an alternative method of cerebrospinal fluid (CSF) diversion used when ventriculoperitoneal or ventriculoatrial shunts are not suitable due to adhesions, a history of peritonitis, ascites, peritoneal dialysis or failure of a prior ventriculoperitoneal shunt.
- Complications of ventriculopleural shunting include pleural effusion, empyema, tension hydrothorax, pneumothorax and rarely a CSF pseudocyst.
- Knowledge of these potential complications is essential when encountering an abnormality of the pleural space in a patient with a ventriculopleural shunt.

Contributors PCT was involved in conception and design, interpretation of imaging, and drafting the article and gave final approval of the version to be published. CS, WJR and TJM (senior author) were involved in analysis of the imaging, revising the article critically for important intellectual content, and gave final approval of the version to be published.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kupeli E, Yilmaz C, Akcay S. Pleural effusion following ventriculopleural shunt: case reports and review of the literature. *Ann Thorac Med* 2010;5:166–70.
- 2 Fox BD, Nayar VV, Johnson KK, et al. Routine imaging in patients with ventriculopleural shunts: lessons learned from a case of tension hydrothorax. *J Neurosurg Pediatr* 2008;2:385.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow