Skin rash from dengue fever

Tsuneaki Kenzaka, Ayako Kumabe

Division of General Medicine,
Center for Community
Medicine, Jichi Medical
University School of Medicine,
Shimotsuke, Tochigi, Japan

Correspondence to

Dr Tsuneaki Kenzaka,
smile.kenzaka@jjichi.ac.jp

DESCRIPTION

A 47-year-old Japanese man presented with a history of having been bitten by mosquitoes, several years earlier in the Philippines, this resulted in a fever and a rash which dissipated spontaneously. He had stayed in the Philippines for 3 weeks, 1 month prior to hospitalisation, and was bitten by several mosquitoes. He had a fever for 6 days and a systemic rash for 3 days before consultation at our hospital. At the time of consultation, his blood pressure was 122/78 mm Hg; pulse 80 bpm, respiratory rate 16 breaths/min and body temperature 38.4°C. Congestion of the palpebral conjunctiva and systemic rash were observed. A tourniquet test on the right arm showed increased petechial haemorrhaging (figure 1). Further, 'islands of white' of the normal skin surrounded by

erythema were noted (figure 2). Blood tests showed the blood platelet count at 52×10^9 cells/L, with abnormal liver function. He received in-patient treatment for suspected dengue fever. A definitive diagnosis of dengue fever was obtained by assessing the levels of dengue virus IgM antibody that was 6.8 (positive cut-off, 1.1 or higher), IgG antibodies 3.5 (positive cut-off, 1.1 or higher) and dengue virus NS1 antigen 18.4 (positive cut-off, 1.0 or higher). A considerable amount of fluid replacement stopped the worsening of his condition and mitigated it, and within 1 week, he was discharged. Even in non-epidemic areas, the diagnosis should be considered in any patient presenting with a characteristic skin rash that has developed within 14 days after even a brief trip to the tropics or subtropics.¹

Figure 1 Tourniquet test positive: showing numerous minute petechial haemorrhages after tourniquet application.

Figure 2 The 'islands of white' of the normal skin surrounded by erythema.

CrossMark

To cite: Kenzaka T,
Kumabe A. *BMJ Case Rep*
Published online: [please
include Day Month Year]
doi:10.1136/bcr-2013-
201598

Learning points

- ▶ Positive tourniquet test and 'islands of white in a sea of red' are characteristic skin rash in dengue fever.
- ▶ Even in non-epidemic areas, the diagnosis should be considered in any patient presenting with characteristic skin rash that has developed within 14 days after even a brief trip to the tropics or subtropics.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCE

- 1 Simmons CP, Farrar JJ, Nguyen VC, *et al.* Dengue. *N Engl J Med* 2012;366:1423–32.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow