Giant left and right atrium: spectrum of rheumatic heart disease

Himanshu Mahla, Shivakumar Bhairappa, Rama Chikkamuniswamy, Cholenahally Nanjappa Manjunath

Department of Cardiology,
Sri Jayadeva Institute of
Cardiovascular Sciences and
Research, Bengaluru,
Karnataka, India

Correspondence to
Dr Shivakumar Bhairappa,
snbhairappa@gmail.com

DESCRIPTION

A 19-year-old man presented with New York Heart Association class II heart failure with a history of rheumatic fever in childhood. He was on regular penicillin prophylaxis, diuretics, vasodilators and β blockers. On examination there was grade IV/VI pansystolic murmur at the apex radiating to the axilla, grade III/VI pansystolic murmur at left sternal border fourth intercostal space (increasing on inspiration). Mean jugular venous pressure was elevated (prominent v waves) with gross pedal oedema. A 12-lead ECG revealed atrial fibrillation with a controlled ventricular rate. A transthoracic echocardiogram revealed severe mitral stenosis (videos 1 and 2), moderate mitral regurgitation (video 3), severe torrential tricuspid regurgitation (video 4) and mild aortic regurgitation (video 5). There was severe right ventricular dysfunction with adequate left ventricular function. Mitral valve

Video 1 Parasternal long axis view showing typical hockey stick mitral valve.

Video 2 Apical 4 chamber view showing severe mitral stenosis with valvar fusion, noncoapting tricuspid valve leaflets and dilated left and right atrium.

Video 3 Apical 4 chamber view with colour doppler across mitral valve showing turbulent forward flow across mitral valve with moderate mitral regurgitation.

Video 4 Apical 4 chamber view with colour doppler across tricuspid valve showing severe tricuspid regurgitation with right ventricular dysfunction.

Video 5 Apical 5 chamber view with colour doppler showing mild aortic regurgitation.

leaflets were thickened and fused with commissural fusion and mild subvalvular disease (figure 1). The mitral valve orifice area was 0.5 cm^2 (figure 1) with gradient of $29/24 \text{ mm Hg}$ (figure 2). The left

To cite: Mahla H, Bhairappa S, Chikkamuniswamy R, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-201853

Figure 1 Basal parasternal short axis view showing mitral valve area.

Figure 2 Apical four-chamber view with continuous wave Doppler across mitral valve with gradient.

atrium was giant with a size of 7.12×8.66 cm (figure 3). The right atrium was also giant with a size of 6.13×8.63 cm (figure 4). Pulmonary pressure was 115 mm Hg (figure 5). The right and left atrial area were 35 and 42 cm² (normal range <20 cm²), respectively. The calculated right and left atrial volumes were 198 and 165 mL (normal range 22–52 mL), respectively. The patient underwent mitral valve replacement, tricuspid valve annuloplasty and left atrial reduction and doing well currently.

This case is a typical example of rheumatic heart disease found in young population in countries like India. Outstanding feature is giant left and right atria both in same patient.

A giant left atrium^{1 2} can be found in 19% of patients requiring surgery for mitral valve. Any left atrial diameter more than 6.5 cm is giant. A giant left atrium (LA) increases the risk of surgery on mitral valve. It is observed that pulmonary oedema occurs less commonly in patients with multiple sclerosis (MS) who have giant

LA. Mitral regurgitation is more commonly associated with giant LA than MS.

A giant right atrium^{3 4} can be due to Ebstein anomaly, pulmonary hypertension or tricuspid regurgitation. In our case severe tricuspid regurgitation with pulmonary hypertension was the cause of the giant RA.

Learning points

- ▶ Rheumatic heart disease is the typical cause of a giant left atrium especially in patients having mitral regurgitation.
- ▶ Patients of rheumatic heart disease with a giant left atrium are protected from pulmonary oedema to some extent.
- ▶ Rheumatic heart disease with a giant right atrium can be seen in patients with severe pulmonary hypertension.

Figure 3 Apical four-chamber view showing left atrial dimensions.

Figure 4 Apical four-chamber view showing right atrial dimensions.

Figure 5 Apical four-chamber view with continuous wave Doppler across tricuspid valve with gradient.

Acknowledgements The authors would like to thank their institute.

Contributors HM and RC diagnosed the case and was involved in the article preparation. SB was involved in corrections. CNM gave guidance and was involved in the final approval of the article.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Apostolakis E, Shuhaiber JH. The surgical management of giant left atrium. *Eur J Cardiothorac Surg* 2008;33:182–90.
- 2 El Maghraby A, Hajar R. Giant left atrium: a review. *Heart Views* 2012;13:46–52.
- 3 Ariyaratnam V, Soni A, Morris A. Giant right atrium in an adult. *Echocardiography* 2008;25:1121–3.
- 4 Gomes S, Wolfenden H, Lambros J. Giant right atrium in an adult: case report of a rare condition. *Heart Lung Circ* 2012;21:50–2.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit

<http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow