Spontaneous haemorrhage in a healthy newborn

Eulália Viveiros,¹ Marcos Sanches,² Alexandra Couto,³ Helena Ramos³

¹Pediatric Department, Hospital do Divino Espírito Santo de Ponta Delgada, EPE, Ponta Delgada, Azores, Portugal

²Pediatric Department, Hospital Distrital de Santarém, EPE, Santarém, Portugal

³Department of Neonatal Intensive Care, Maternidade Dr Alfredo da Costa, Lisboa, Portugal

Correspondence to

Dr Eulália Viveiros,
eulaliaviveiros@gmail.com

Accepted 3 May 2014

DESCRIPTION

A full-term male neonate, who was born by vaginal delivery without trauma, presented a small purple tumefaction on his back at 8 h of life, with a tender consistency. It was well defined and interpreted as a haemangioma. At 24 h of life he became lethargic, pale, tachycardic and the tumefaction worsened (figures 1 and 2).^{1 2} A careful review of his family history revealed haemophilia A in first-degree to third-degree relatives on the maternal side.^{1 2} Haemoglobin dropped from 14.6 to 9.5 g/dL in 4 h. His activated thromboplastin time was 85.2 s with normal prothrombin time and number of platelets. The coagulation factor dosage revealed a severe factor VIII deficiency (FVIII <0.1%). He received red blood cells, fresh frozen plasma and intravenous recombinant factor VIII transfusions, and started phototherapy for jaundice.² Cranial, abdominal ultrasounds and thorax radiographies were normal. In the following days, his general condition and haematoma significantly improved.

He was discharged at 7 days of life with FVIII dosage in normal range (85.2%) and negative inhibitors for FVIII.

Learning points

- ▶ Haemophilia A is an X-linked coagulation disorder, so mothers are usually asymptomatic carriers and manifestations are almost exclusively in male neonates.
- ▶ A review of the family history in an otherwise healthy newborn, with unexplained bleeding, can lead to the diagnosis in two-thirds of the population.
- ▶ The severe form of haemophilia should be part of the differential diagnosis of haemodynamically unstable newborns with spontaneous superficial haematoma.

Figure 1 Diffuse haematoma on the neonate's back and the presence of some central yellowish papule surrounded by a halo of erythema (erythema toxicum).

CrossMark

To cite: Viveiros E, Sanches M, Couto A, *et al.* *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-202253

Figure 2 Extensive haematoma on the back of the neonate.

Acknowledgements The authors would like to thank all in the neonatal intensive care and haematology department.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Chalmers EA. Neonatal coagulation problems. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F475–8.
- 2 Dyson H. Neonatal haemophilia—a guide for recognition and management. *Haemophilia* 2006;2:156–9.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow