Drop attacks: an unexpected diagnosis

Ana Cristina Lladó, Rui Malheiro, Ana Rita Estriga

Department of Internal Medicine, Hospital Santo António dos Capuchos, Lisbon, Portugal

Correspondence to

Dr Ana Cristina Lladó, ana.llado@sapo.pt

Accepted 19 February 2014

DESCRIPTION

A healthy 40-year-old woman, without previous medication presented with 6-month history of recurrent falls, with no prodromes or triggering factor, loss of consciousness and with instantaneous recovery to baseline status. Physical examination, blood tests, ECG and echocardiography, EEG were unrevealing. CT of the brain showed a giant cystic lesion of the posterior fossa with mass effect and shaping of fourth ventricle and cisterns of the base (figure 1A). MRI revealed a lesion with an identical signal to the cerebrospinal fluid with no abnormal enhancement after gadolinium injection, suggesting invaginated arachnoid cyst (figure 1B,C). Patient underwent craniotomy and cyst fenestration to the magna cistern (figure 2). It was a successful surgery which permitted a gain in quality of life because no further falls were seen.

Drop attacks consist of sudden falls like described above. This clinical picture is more frequent in elderly individuals and idiopathic in 65% of cases. It is usually explained by a transient bilateral disorder involving structures of the central nervous system responsible for postural tonus and balance. However, over time the term 'drop attacks' was applied to different situations associated with fall without meeting the defining criteria noted above (syncope, arrhythmias, vasovagal responses, transient ischaemia of the vertebrobasilar area, epileptic variants, Tumarkin otolithic crises, psychiatric disorders, etc). These ones must be assumed as possible differential diagnoses.¹⁻³ The treatment must be individualised. In this case we have a middle-age patient with a median lesion of posterior fossa implicated in a transient blocking of the circulatory cerebral aqueduct.

Figure 2 Control CT after surgery showed a persistent but smaller lesion with a considerable reduction of mass effect.

Learning points

- ▶ Drop attacks consist of sudden falls without prodromes, trigger factors or loss of consciousness and with instantaneous recovery to baseline status.
- ▶ They are usually explained by a transient bilateral disorder involving structures of the central nervous system responsible for postural tonus and balance.
- ▶ Several differential diagnoses should be considered.

Figure 1 CT of the brain showed a giant cystic lesion of the posterior fossa with mass effect and shaping of fourth ventricle and basal cisterns (A—axial view). MRI revealed lesion with an identical signal to cerebrospinal fluid (B—axial view and C—sagittal view) with no abnormal reinforcements after gadolinium injection, suggesting invaginated arachnoid.

CrossMark

To cite: Lladó AC, Malheiro R, Estriga AR. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-202551

Contributors All authors contributed to the clinical evaluation of the patient and to the diagnostic approach. They were also involved in the bibliographic research and image collection.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Lanska DJ. Drop attacks. *Neurology* 2004 [medLink]. retrieved 18 October 2013. <http://www.medlink.com/medlinkcontent.asp>
- 2 Welsh LW, Welsh JJ, Lewin B, *et al.* Vascular analysis of individuals with drop attacks. *Ann Otol Rhinol Laryngol* 2004;113:245.
- 3 Dallan I, Bruschini L, Nacci A. Drop attacks and vertical vertigo after transtympanic gentamicin: diagnosis and management. *Acta Otorhinolaryngol Ital* 2005;25:370.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow