Ladakhi koilonychia

Uday Yanamandra,¹ Ruchira Mukherji,¹ Sagarika Patyal,¹ Velu Nair²

¹Department of Ophthalmology, 153 General Hospital, Leh, Jammu and Kashmir, India
²Armed Forces Medical College, Pune, Maharashtra, India

Correspondence to
 Dr Uday Yanamandra,
 udayj2@gmail.com

DESCRIPTION

A 48-year-old native highlander of the Leh Ladakh region (approximate height 12 000ft) was diagnosed with severe koilonychia after visiting a medical camp (figures 1–4).

On examination, his built was normal, there was no pallor and no other signs of nutritional deficiency. Abdominal, respiratory, neurological and cardiovascular examinations were normal. The patient's haemoglobin was 16.1 g/dL. All other haematological and biochemical parameters were normal. Peripheral blood smear revealed a normal picture. Iron studies revealed serum iron of 105 (65–175) µg/dL, the total iron binding capacity was 259 (250–450) µg/dL, transferrin saturation was 18.2 (13–45)% and serum ferritin 166.7 (22–322) µg/dL.

Important differentials for a patient with koilonychia include chronic iron deficiency, trauma,

haemochromatosis, thyroid dysfunction, high altitude, exposure to petrochemicals and Plummer-Vinson syndrome.^{1 2} Detailed history and examination ruled out the above causes except the exposure to high altitude as a cause for koilonychia in our patient.

Exposure to high altitude is a known aetiology for koilonychias, also described by some authors as 'Ladakhi koilonychia'.³ The pathophysiology of the commonly found koilonychias in native highlanders is still not well understood; though some describe it to be seasonal, secondary to anaemia or covert iron deficiency due to increased erythropoiesis.³ Our patient had none of these described pathophysiological elements.

This image demonstrates the likelihood of development of koilonychia in a well-adapted native highlander.

Figure 1 Images of Ladakhi koilonychia in frontal view.

Figure 3 Spooned nails with oil drops retained by nails in frontal view.

Figure 2 Images of Ladakhi koilonychia in profile.

Figure 4 Spooned nails with oil drops retained by nails in profile.

To cite: Yanamandra U, Mukherji R, Patyal S, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-202567

Learning points

- ▶ High altitude exposure is a known cause of koilonychia, described by some as Ladakhi koilonychia.
- ▶ The aetiology of this koilonychia needs to be ascertained and is not always a consequence of iron deficiency.
- ▶ Before labelling high altitude as a cause of koilonychia, all other treatable causes of koilonychia should be ruled out.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Fawcett RS, Linford S, Stulberg DL. Nail abnormalities: clues to systemic disease. *Am Fam Physician* 2004;69:1417–24.
- 2 Novacek G. Plummer-Vinson syndrome. *Orphanet J Rare Dis* 2006;1:36.
- 3 Sawhney MP. Ladakhi koilonychia. *Indian J Dermatol Venereol Leprol* 2003;69:79–80.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow