Rapid diagnosis of *Campylobacter jejuni* by stool Gram stain examination

Kazuhiro Kamata,¹ Yasuharu Tokuda²

¹Mito Kyodo General Hospital, Mito City, Japan

²Department of General Internal Medicine, Tsukuba University, Mito, Ibaraki, Japan

Correspondence to

Professor Yasuharu Tokuda, yasuharu.tokuda@gmail.com

DESCRIPTION

Microscopic stool examination by Gram stain is useful for rapid diagnosis and treatment of *Campylobacter* infection in patients with acute onset diarrhoea. A 69-year-old Japanese woman with pernicious anaemia presented with a 3-day history of fever and watery diarrhoea. On examination, her mental status was normal and general appearance was not in acute distress. Blood pressure was 115/43 mm Hg, pulse rate 131 bpm and regular, respiration rate 20 breaths/min and temperature 39°C. The palpebral conjunctivae were anaemic and tongue surface was atrophic and dry. Cardiovascular and lung examinations revealed no abnormality except for tachycardia. Abdominal examination revealed normal bowel sounds and mild tenderness on the right lower quadrant without rebound tenderness. Laboratory data showed elevated leucocyte count and macrocytic anaemia. The faecal Gram stain examination revealed numerous Gram-negative rods of gull-winged form (figure 1). Oral erythromycin was prescribed based on a diagnosis of *Campylobacter*

enteritis. Two days after the first visit, her symptoms were improved and the stool culture grew *Campylobacter jejuni*. *Campylobacter* species are a frequent cause of infectious enteritis. Although its definitive diagnosis requires culture results, the stool Gram stain finding of gull-wing-shaped Gram-negative rods has good sensitivity and specificity.^{1 2} The Gram stain examination has low cost and provides real-time diagnosis that can lead to immediate and appropriate treatment.

Learning points

- ▶ Stool Gram stain examination leads to timely diagnosis and treatment of *Campylobacter* infection in patients with acute onset diarrhoea.
- ▶ Gram stain examination is a precious and inexpensive tool for managing common infectious diseases.
- ▶ The presence of Gram negative rods with gull-wing shape in stools of diarrhoea patients suggests *Campylobacter jejuni* enteritis.

Figure 1 White arrow indicates gull-wing-shaped Gram-negative rods.

Contributors YT cared the patient and YT and KK wrote the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Park CH, Hixon DL, Polhemus AS, *et al*. A rapid diagnosis of *Campylobacter* enteritis by direct smear examination. *Am J Clin Pathol* 1983;80:388–90.
- 2 Wang H, Murdoch DR. Detection of *Campylobacter* species in faecal samples by direct Gram stain microscopy. *Pathology* 2004;36:343–4.

CrossMark

To cite: Kamata K, Tokuda Y. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-202876

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>. BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow