CT imaging is invaluable in diagnosing emphysematous pyelonephritis (EPN): a rare urological emergency

Ali Moostapha Tasleem,¹ Paul Murray,² Faqar Anjum,¹ Seshadri Sriprasad¹

¹Department of Urology, Darent Valley Hospital, Kent, UK
²Darent Valley Hospital, Kent, UK

Correspondence to
 Ali Moostapha Tasleem,
 amt03@doctors.org.uk

Accepted 16 March 2014

DESCRIPTION

Emphysematous pyelonephritis (EPN) is an acute, fulminant and potentially fatal necrotising process with varying clinical presentations and a radiological classification. Most cases occur in diabetics with poor glycaemic control, while a minority are due to urinary tract obstruction. Early CT imaging is vital in diagnosing highly morbid EPN, with mortality rates of up to 69% described, and can necessitate nephrectomy.^{1 2}

A 79-year-old woman, with non-insulin-dependent diabetes mellitus presented with confusion, fever, rigours and diffuse abdominal pain. Laboratory tests revealed a neutrophilia predominant leucocytosis,

Figure 1 Axial CT section showing gas within the collecting system of the left kidney and perinephric stranding.

Figure 2 Sagittal CT section displaying an impressive volume of gas and resulting air-fluid level in the left kidney.

Figure 3 Sagittal CT section showing resolution of the gas within the renal collecting system and a decrease in inflammation. This corresponded to an improvement in the patient's clinical state.

normal renal function and metabolic acidosis on arterial analysis. Prompt contrast-enhanced CT imaging clinched the diagnosis of left type I EPN, revealing an obstructing ureteric calculus and demonstrating air-fluid levels with an impressive volume of gas in the collecting system (figures 1 and 2). Vigorous resuscitation, inotropic support and the insertion of an ultrasound-guided nephrostomy with broad-spectrum antibiotics given initially formed the cornerstone of management. Urine and blood cultures returned positive for *Escherichia coli* and intravenous amikacin was administered for 10 days. She remained an inpatient for 4 weeks requiring a programme of supplementary nasogastric feeding and a structured rehabilitation programme of intensive physiotherapy. She gradually improved clinically and follow-up CT imaging displayed resolution of the gas and a decrease in the associated perinephric stranding and infection (figure 3).

Learning points

- ▶ Suspect emphysematous pyelonephritis (EPN) in unwell elderly patients with diabetes with a fever of unknown origin, or whose clinical presentation is suggestive of a urinary tract infection.
- ▶ Prompt CT imaging is vital in establishing a diagnosis of EPN and ruling out other causes for intra-abdominal sepsis.
- ▶ Early and aggressive resuscitation, empirical antimicrobial therapy and a drainage nephrostomy tube are imperative for a positive clinical outcome.

To cite: Tasleem AM, Murray P, Anjum F, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-204040

Our case demonstrates the high index of clinical suspicion required to diagnose EPN and aggressive subsequent treatment strategies. We advocate a multidisciplinary approach with prompt CT imaging, which is instrumental in categorising the severity and type of EPN, thus guiding management.³

Acknowledgements The authors would like to thank the medical records department.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Saxena D, Aggarwal L, Tudu SK, *et al*. Emphysematous pyelonephritis—a rare surgical emergency presenting to the physician: a case report and literature review. *Indian J Surg* 2013;75 (Suppl 1):272–4.
- 2 Huang JJ, Tseng CC. Emphysematous pyelonephritis: clinoradiological classification, management, prognosis, and pathogenesis. *Arch Intern Med* 2000;160:797–805.
- 3 Pontin AR, Barnes RD. Current management of emphysematous pyelonephritis. *Nat Rev Urol* 2009;6:272–9.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow