Gregg syndrome aka embryopathia rubeolaris: CT illustration

Ajit Kumar Reddy,¹ Sathya Narayanan Renganathan,² Annitha Elavarasi Jayamohan,¹ Prakash Manikka Lakshmanan²

¹Mahatma Gandhi Medical College and Research Institute, Pondicherry, India

²Department of Radiology, Mahatma Gandhi Medical College and Research Institute, Pondicherry, India

Correspondence to
Dr Ajit Kumar Reddy,
drajitreddy@gmail.com

Accepted 25 May 2014

DESCRIPTION

Introduction

Congenital rubella syndrome (CRS) is a spectrum of clinical findings in a patient that results from primary maternal infection with rubella virus. Diagnosis is usually made clinically but imaging findings can prompt the diagnosis in appropriate clinical setting. The features of congenital rubella include sensorineural deafness, cataract, cardiac anomalies, mental retardation and microcephaly.

Case report

We report a case of 10-year-old girl who presented with a history of global developmental delay, intellectual disability and having been treated for patent

ductus arteriosus. There was no maternal history of exposure to infection or vaccination during her pregnancy. The girl was the second child of a non-consanguineous marriage with no similar symptoms in her elder sibling. Clinical examination revealed mild facial asymmetry and hearing loss with bilateral central cataract (R>L; [figure 1](#)). CT of the brain ([figure 2](#)) showed bilateral dense, coarse calcifications of basal ganglia and a linear calcification was also noted in the right frontal lobe in the subcortical white matter.

Differential diagnosis

Basal ganglia and cortical calcifications are common features of all infections that constitute the toxoplasmosis, rubella, cytomegalovirus, herpes simplex virus (TORCH) syndrome. Cytomegalovirus and toxoplasmosis cause bilateral periventricular and subependymal calcifications.¹ Interestingly, calcifications of toxoplasmosis may be resolved with therapy.² Calcification pattern of TORCH infection is not specific for any infection.^{2,3}

Endocrinal disorders such as hypoparathyroidism, pseudohypoparathyroidism, hypothyroidism and metabolic causes such as Leigh's disease, Fahr's disease, Cockayne's syndrome are well-known causes of basal ganglia calcifications.

Figure 1 Clinical photograph of a 10-year-old girl shows bilateral cataract (right>left).

Figure 2 CT of the brain reveals bilateral dense basal ganglia calcification (block arrow) and subcortical white matter calcification in the right frontal lobe (thin arrow).

To cite: Reddy AK, Renganathan SN, Jayamohan AE, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-204204

Similarly carbon monoxide, lead poisoning and postchemotherapy/radiation therapy result in calcifications as well. However a normal variant of basal ganglia calcification is not uncommon.

Among the above causes cataract may be seen in toxoplasmosis, hypoparathyroidism, hypothyroidism, Cockayne's syndrome and postradiation therapy. However, the cataracts in CRS are often bilateral, central and may be lamellar, nuclear or membranous in nature.

Learning points

- ▶ Classical triad of congenital rubella syndrome is sensorineural deafness, cataract and cardiac anomalies.
- ▶ Clinical manifestations of rubella are more severe if infection occurs in early trimester.
- ▶ Bilateral periventricular, basal ganglia and brainstem calcifications are found.

Contributors AKR acquired data, drafted the article, analysed and interpreted the data, revised the article critically for important intellectual content and finally approved the version for publishing. SN acquired data, drafted the article and finally approved the version for publishing. AEJ analysed and interpreted data, revised the article critically for important intellectual content, finally approved the version for publishing. MLP analysed and interpreted data, revised the article critically for important intellectual content and finally approved the version for publishing.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 KiroUglu Y, Ccaltl C, Karabulut N, *et al*. Intracranial calcifications on CT. *Diagn Interv Radiol* 2010;16:263–9.
- 2 Appliedradiology.com. Articles: intracranial calcifications on applied radiology. <http://www.appliedradiology.com/Issues/2009/11/Articles/Intracranial-calcifications/Intracranial-calcifications.aspx> (accessed 6 Feb 2014).
- 3 Osborn A. *Osborn's brain*. Salt Lake City, UT: Amirsys Pub, 2013.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow