Interstitial lung disease in antisynthetase syndrome without clinical myositis

Pranathi Rao Sundaragiri,¹ Saraschandra Vallabhajosyula,¹ Jennifer Papa Kanaan²

¹Department of Internal Medicine, Alegant-Creighton University Medical Center, Creighton University School of Medicine, Omaha, NE, USA

²Division of Pulmonary, Critical Care and Sleep Medicine, Department of Internal Medicine, University of Connecticut Health Center, Farmington, CT, USA

Correspondence to

Dr Saraschandra Vallabhajosyula, SaraschandraVallabhajosyula1@creighton.edu

Accepted 11 March 2014

DESCRIPTION

Antisynthetase syndrome (ASS) is a lesser recognised idiopathic inflammatory myopathy with nearly 89% showing interstitial lung disease (ILD).¹ However, in a small subpopulation without evidence of myositis, the diagnosis may be critically delayed, hindering management of this rapidly progressive disease.

A 58-year-old Caucasian man presented with polyarthritis involving bilateral knees, shoulders, wrists and hand joints. He reported a history of questionable asthma with progressive dyspnoea despite optimal therapy. Examination revealed coarse pulmonary crepitations and diffuse small-joint swelling with rhagadiform skin changes over hands (figure 1). Laboratory studies demonstrated leucocytosis and modest creatinine kinase elevation. Pulmonary function tests showed forced expiratory volume in 1 s/forced vital capacity (FEV₁/FVC) 76%, diffusing capacity of the lungs for carbon monoxide (DLCO) 63%, FEV₁ 1.99 L (54%) with 9% (230 mL) reversibility with β -agonists, consistent with restrictive lung disease. High-resolution CT of the chest noted diffuse septal thickening and honeycombing (figures 2 and 3). Inflammatory arthritis, hyperkeratotic 'mechanic' hands and ILD prompted a diagnosis of ASS, confirmed by serum anti-Jo1 antibodies, despite the absence of clinical myositis. The patient improved on prednisone and methotrexate, and was later transitioned to mycophenolate mofetil.

Only 2–11% patients with ASS present with ILD in the absence of myositis and are diagnosed with anti-Jo1 antibodies in the serum.² Median interval

Figure 2 High-resolution CT of the chest demonstrating honeycombing and septal thickening consistent with interstitial lung disease.

of development of myositis after ILD is 6 months; however rarely extending up to 2 years.^{1–3} Anti-Jo1 antibody-positive ILDs cause only modest serum creatine phosphokinase elevation that delays diagnosis in the absence of clinical myositis.¹ These patients deteriorate faster in comparison to patients with myositis. However, once diagnosed, they have high steroid-responsiveness, which is postulated to be due to higher non-specific interstitial pneumonia without manifest alveolar damage.² High-resolution CT scan has replaced lung biopsy in establishing diagnosis. Steroid-resistant cases are treated with azathioprine or cyclophosphamide.^{1,2}

Figure 1 Hyperkeratotic 'mechanic' hands.

Figure 3 Prominent interstitial thickening consistent with interstitial lung disease.

CrossMark

To cite: Sundaragiri PR, Vallabhajosyula S, Kanaan JP. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-204296

Learning points

- ▶ Clinical features and diagnosis of antisynthetase syndrome (ASS) including interstitial lung disease and 'mechanic' hands.
- ▶ Recognising lesser known manifestations of ASS in the absence of clinical myositis.

Contributors PRS was involved in clinical evaluation, literature search, drafting and finalising the manuscript. SV was involved in literature search, critical analysis,

drafting and finalising the manuscript. JPK was involved in clinical evaluation, critical analysis and finalising the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Marie I, Josse S, Hatron PY, *et al.* Interstitial lung disease in anti-Jo1 patients with antisynthetase syndrome. *Arthritis Care Res (Hoboken)* 2013;65:800–8.
- 2 Plastiras SC, Soliotis FC, Vlachoyiannopoulos P, *et al.* Interstitial lung disease in a patient with antisynthetase syndrome and no myositis. *Clin Rheumatol* 2007;26:108–11.
- 3 Patel HC, Lauder NN. The antisynthetase syndrome. *Am J Med* 2011;124(9):e3–4.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow