CASE REPORT

Sturge-Weber syndrome: oral and extra-oral manifestations

Amitandra Kumar Tripathi,¹ Vivek Kumar,² Rahul Dwivedi,³ Charanjit Singh Saimbi⁴

¹Department of Periodontology, Career Postgraduate Institute of Dental Sciences and Hospital, Lucknow, Uttar Pradesh, India

²Department of Periodontology, Career Postgraduate Institute of Dental Sciences and Hospital, Lucknow, Uttar Pradesh, India

³Department of Pedodontics, Career Postgraduate Institute of Dental Sciences and Hospital, Lucknow, Uttar Pradesh, India

⁴Department of Periodontology, King George's Medical University, Lucknow, Uttar Pradesh, India

Correspondence to
Professor Charanjit Singh Saimbi, cssaimbi@gmail.com

Accepted 19 February 2015

SUMMARY

Sturge-Weber syndrome is a rare, congenital, neuro-oculo-cutaneous disorder which is characterised extra-orally by unilateral port wine stains on the face, glaucoma, seizures and mental retardation, and intra-orally by ipsilateral gingival haemangioma which frequently affects the maxilla or mandible. In the present case, a 15-year-old female patient presented with a port wine stain on the right side of the face and glaucoma of the right eye, and intra-orally with gingival haemangioma on the right side of the maxilla.

BACKGROUND

Sturge-Weber syndrome (SWS) or encephalotrigeminal angiomatosis belongs to a group of disorders known as phakomatoses. In SWS, abnormal development of the primordial vascular bed occurs during the early stages of cerebral vascularisation, when the blood supply to the brain, meninges and face is undergoing reorganisation. As a result of these abnormal changes, the overlying leptomeninges are richly vascularised and the brain beneath becomes atrophic and calcified, mainly in the molecular layer of the cortex.

The main characteristic features of this disorder are port wine stains on the forehead and/or scalp, seizures and mental retardation due to angioma formation in the brain, and glaucoma.¹

Schirmer provided the first detailed description of SWS in 1860, Sturge further described SWS-related dermatological, ophthalmic and neurological manifestations in 1879, and Weber reported radiological alterations seen in these patients in 1929.² This disorder occurs in 1:50 000 live births.³

Oral manifestations include unilateral haemangiomatous lesions in the maxillary or mandibular gingiva, lips, tongue, palatine region, etc. In gingiva, these lesions present as unilateral hyperplasia due to an increase in vascular component, and demonstrate bleeding following minimal trauma.¹

CASE PRESENTATION

A 15-year-old female patient presented to the periodontology department with a 3–4-month history of pain and bleeding from the gums while eating. The patient's past medical history revealed that she had developed a convulsive disorder at the age of 1 year for which she was taking medication.

Extra-oral examination showed a port wine stain on the right side of the face (figure 1), while eye examination showed dilated blood vessels in the right eye (figure 2). Intra-oral examination showed

Figure 1 Port wine stain on the right side of the face.

unilateral hyperplastic lesions on the right side of the maxilla (figure 3A, B). These lesions showed blanching when pressure was applied (figure 4). The patient had poor oral hygiene.

On radiographic examination, a lateral cephalogram and postero-anterior view showed tram-line calcification (figures 5 and 6). CT showed severe alveolar bone loss on the affected side of the oral cavity (figure 7). Blood investigations were normal. Based on the history, and on clinical and radiological findings, a diagnosis of SWS was made.

The patient was treated with oral prophylaxis, advised to use a chlorhexidine mouth rinse and

Figure 2 Dilated blood vessels in the right eye.

CrossMark

To cite: Tripathi AK, Kumar V, Dwivedi R, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-207663

Figure 3 (A, B) Unilateral hyperplastic lesions on the right side of the maxilla.

given oral hygiene instructions. The patient and her parents were referred for psychological counselling.

INVESTIGATIONS

On radiographic examination, a lateral cephalogram and postero-anterior view showed tram-line calcification. CT showed severe alveolar bone loss on the affected side of the oral cavity. Blood investigations were normal.

Figure 4 Blanching of lesion under pressure.

Figure 5 Lateral cephalogram showing tram-line calcification in the occipital region.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis included Rendu-Osler-Weber syndrome, angio-osteodystrophy syndrome, Maffucci syndrome and Klippel-Trenaunay-Weber syndrome.

TREATMENT

Treatment consisted of establishing an oral plaque control regimen for the patient. Oral prophylaxis (scaling and root planing) was carried out and the patient was instructed on oral hygiene and the use of chlorhexidine mouth rinse. The patient and her parents were referred for psychological counselling.

Figure 6 Postero-anterior view showing tram-line calcification in the parietal region.

Figure 7 CT showing severe alveolar bone loss on the affected side of the oral cavity.

OUTCOME AND FOLLOW-UP

At the 1-month and 3-month follow-up visits, there was a significant improvement in the patient's chief complaint and maintenance of oral hygiene was noted.

DISCUSSION

SWS consists of a group of disorders which may present with neurological, cutaneous, ocular and oral manifestations. The most common clinical characteristic is a port wine stain on the face which normally follows the course of the trigeminal nerve, especially the ophthalmic division.⁴ Port wine stains are reported in about 87–90% of cases and generally occur on the right side, although these lesions present bilaterally in about 33% of cases.⁵ In the present case, the port wine stain was present on the right side of the face.

Leptomenigeal angiomatosis is another important clinical feature of this syndrome, and can lead to contra-lateral hypertrophy, progressive cerebral calcification, epileptic convulsions and mental retardation.⁴ In this case, the patient had a history of epileptic convulsions and her skull radiograph showed intracranial calcification in the occipito-parietal region.

According to the Roach scale, SWS is classified as follows:⁶

Type I: both facial and leptomenigeal angiomatosis; may have glaucoma

Type II: facial angiomatosis only (no CNS involvement); may have glaucoma

Type III: isolated leptomenigeal angiomatosis; usually no glaucoma.

According to the above criteria, our case was type I SWS as the patient had a port wine stain and glaucoma on the right side with epileptic convulsions and calcification of the occipito-parietal region.

The most common oral manifestations in SWS are mainly seen in the unilateral side of the gingiva and finish abruptly at the midline. Angiomatous gingival lesions range from slight vascular hyperplasia to severe haemangiomatous proliferation due to abnormal increases in the vascular component. As a result of these changes, gingiva bleed on slight provocation or minor trauma.²

In the present case, massive haemangiomatous proliferation was seen on the right side of the maxillary gingiva which blanched on the application of pressure and bled profusely following slight trauma.

Learning points

- ▶ Sturge-Weber syndrome presents with a large number of clinical manifestations and early diagnosis is difficult.
- ▶ Early diagnosis is necessary to allow future complications, mainly CNS related, to be controlled.
- ▶ Expert psychological counselling of patients and their parents is necessary.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Royle HE, Lapp R, Ferrara ED. The Sturge-Weber syndrome. *Oral Surg Oral Med Oral Pathol* 1966;22:490–7.
- 2 Neto FXP, Junior MAV, Ximenes LS, et al. Clinical features of Sturge-Weber syndrome. *Int J Arch Otorhinolaryngol* 2008;12:565–70.
- 3 Welty LD. Sturge-Weber syndrome: a case study. *Neonatal Netw* 2006;25:89–98.
- 4 Mukhopadhyay S. Sturge-Weber syndrome: a case report. *J Indian Soc Pedod Prev Dent* 2008;26:29–30.
- 5 Inan CJM. Sturge-Weber syndrome; report of an unusual cutaneous distribution. *Brain Develop* 1999;21:68–70.
- 6 Del Monte M, Eisbschitz-Tsimhoni M. Sturge-Weber syndrome: overview. *eMedicine* February, 2007.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow