Simple leads to complex solutions

Mathew Sheridan,¹ Senthil Kirubakaran²

¹Eastbourne District General Hospital, Eastbourne, East Sussex, UK

²Queen Alexandra Hospital, Portsmouth, UK

Correspondence to

Dr Mathew Sheridan,
mda04mjs@gmail.com

Accepted 14 January 2015

DESCRIPTION

A 76-year-old woman on warfarin for atrial fibrillation presented for routine pacing check 6 weeks after implantation of a dual-chamber device. She felt weak and generally unwell. The atrial lead was not sensing or pacing at high outputs. A chest X-ray showed the lead had perforated the atrial wall (figure 1). Echocardiography showed no pericardial effusion. International normalised ratio was 2.4.

Figure 1 Chest radiograph 6 weeks after device implantation showing the atrial lead has perforated the atrial wall.

In light of these findings, urgent lead revision was planned in a centre with cardiothoracic facilities due to the increased risk of pericardial effusion and tamponade.

Although pacemaker malfunction can be complex, simple non-specialist investigations can elucidate causes and majorly impact management.

Learning points

- ▶ Simple, systematic investigation of unwell patients (with cardiac devices) before specialist input can majorly inform and impact management.
- ▶ Delayed lead perforation is defined as lead perforation more than 30 days after implantation and has a reported incidence of 0.8%.¹
- ▶ Although many cases of lead perforation may be asymptomatic,¹ prompt recognition (facilitated by simple investigation) may prevent resultant pericardial effusion, tamponade and fistulae.¹

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCE

- 1 Khoeiry G, Lakhani M, Abi Rafeh N, *et al*. Right coronary artery fistula as a result of delayed right atrial perforation by a passive fixation lead. *Circ Arrhythm Electrophysiol* 2012;5:e46–7.

CrossMark

To cite: Sheridan M, Kirubakaran S. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-208869

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow