Inflammation of actinic keratoses during paclitaxel chemotherapy

Faisal R Ali, Zenas Z N Yiu, David Fitzgerald

Department of Dermatology,
Salford Royal NHS Foundation
Trust, Salford, UK

Correspondence to

Dr Zenas Zee Ngai Yiu,
zenas.yiu@manchester.ac.uk

Accepted 11 May 2015

DESCRIPTION

A 73-year-old woman with a long-standing history of sun exposure was being treated for breast cancer with paclitaxel chemotherapy. Following the third cycle of chemotherapy, she developed an inflamed, 'sore' rash on her chest. Examination revealed numerous inflamed, hyperkeratotic papules superimposed on an erythematous background on the upper back, anterior upper chest and forearms, in a striking photodistributed pattern (figure 1).

The patient had multiple actinic keratoses (AKs) prior to starting chemotherapy, which became inflamed following instigation of systemic paclitaxel. AKs are the commonest premalignant skin pathology with the potential to progress to squamous cell carcinoma. Inflammatory changes are often observed in patients with AKs taking other chemotherapeutic agents, most notably 5-fluorouracil and capecitabine.^{1 2} Following initiation of chemotherapy, patients develop inflammation and pruritus of the AKs, which, if left untreated, continue throughout the duration of chemotherapy and subside following completion of treatment, with apparent resolution of the previously inflamed AKs. The chemotherapy is

presumed to selectively induce DNA damage within sun-damaged keratinocytes.³

Physicians should be aware of this potential side-effect of chemotherapeutic agents in patients with previously high sun exposure. Such cutaneous manifestations do not represent an allergic reaction requiring cessation of life-saving chemotherapy, but, rather, indicate elimination of underlying AKs, and can be successfully treated with topical corticosteroids.

Learning points

- ▶ Actinic keratoses are the commonest premalignant skin lesion with the potential to progress to squamous cell carcinoma.
- ▶ Pre-existing actinic keratoses (typically on sun-exposed sites) may become inflamed following instigation of chemotherapy.
- ▶ This inflammatory response does not necessarily require discontinuation of the chemotherapy, but can be treated with topical corticosteroids.

Contributors DF, FA and ZZNY were responsible for drafting the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Johnson TM, Rapini RP, Duvic M. Inflammation of actinic keratoses. *J Am Acad Dermatol* 1987;17:192–7.
- 2 Krathen M, Treat J, James WD. Capecitabine induced inflammation of actinic keratoses. *Dermatol Online J* 2013;13:13.
- 3 Fu W, Cockerell CJ. The actinic (solar) keratosis: a 21st-century perspective. *Arch Dermatol* 2003;139:66–70.

Figure 1 Inflammation of actinic keratoses following instigation of paclitaxel chemotherapy.

CrossMark

To cite: Ali FR, Yiu ZN, Fitzgerald D. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-209925

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow