Assessment of severe lower limb trauma with the aid of on-site photography

Conor Gouk,¹ Narayan Mohanakrishnan,² Emma Jennifer Keen³

¹Department of Orthopaedics, Gold Coast University Hospital, Gold Coast, Queensland, Australia

²Greenslopes Private Hospital, Brisbane, Queensland, Australia

³Gold Coast University Hospital, Gold Coast, Queensland, Australia

Correspondence to

Dr Conor Gouk, c.j.gouk.06@aberdeen.ac.uk

Accepted 19 March 2015

DESCRIPTION

A 44-year-old man was brought to the hospital by ambulance with a severely deformed right femur following a motorbike accident. The initial deformity was reduced on-site, but the ambulance crew provided a pre-reduction photograph (figure 1). The patient was travelling at 100 kmph when he collided with a wall. Evaluation on-site, and primary and secondary surveys, revealed that the patient was stable (as per Advanced Trauma Life Support (ATLS) protocol) and his injured limb was neurovascularly intact.

Owing to the severity of his initial deformity, a CT angiogram was performed. Examination and imaging revealed a compound, comminuted right femoral fracture involving the distal third of the femoral shaft and femoral condyles with associated fracture of the patella (figures 2 and 3). However, there was no evidence of arterial injury. No other injuries were identified.

Figure 1 On-site photograph of the injured lower limb.

Emergency open reduction and internal fixation was performed using headless compression screws and a lateral locking plate (figure 4). Postoperatively, the patient was placed in a range of motion brace

Figure 2 Plain anteroposterior radiograph of the distal femur.

Figure 3 CT slice of the distal femur showing the extent of the injury.

CrossMark

To cite: Gouk C, Mohanakrishnan N, Keen EJ. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-210155

Figure 4 Plain anteroposterior radiograph of the fracture post open reduction internal fixation with accompanying range of motion brace.

locked in extension, and was non-weightbearing for 6 weeks. He was discharged from hospital 5 days postoperatively without complication.

High-energy blunt trauma to the lower extremity is associated with a 28–46% rate of injury to the popliteal artery.¹ Delay in diagnosis and failure to revascularise within the first 6–8 h lead to high rates of limb amputations.² We present this as an interesting case of severe limb injury, with on-site photography used to guide emergency teams in assessment, management and consideration of concomitant injury.

Learning points

- Importance of the first responder handover and the potential aid photographic adjuncts offer when conveying vital information in handover to emergency staff.
- Importance of neurovascular assessment of patients with trauma.
- Importance of radiological adjunct imaging in trauma settings.

Acknowledgements The authors wish to thank Dr Jason Beer, the emergency staff at Gold Coast University Hospital and the Queensland Ambulance Service.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Chapman JA. Popliteal artery damage in closed injuries of the knee. *J Bone Joint Surg Br* 1985;67:420–3.
- 2 Steele HL, Singh A. Vascular injury after occult knee dislocation presenting as compartment syndrome. *J Emerg Med* 2012;42:271–4.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow