Leptomeningeal metastases in pineoblastoma

Sonia Sandip,¹ Chandan J Das,² Rohit Kumar Khandelwal³

¹All India Institute of Medical Sciences, New Delhi, India
²Department of Radiodiagnosis, All India Institute of Medical Sciences, New Delhi, India
³Department of Radiology, All India Medical Sciences, New Delhi, India

Correspondence to
 Dr Chandan J Das,
 dascj@yahoo.com

Accepted 6 June 2015

DESCRIPTION

Pineoblastoma is one of the pineal parenchymal tumours, which are rare tumours and account for approximately 15% of all tumours found in the pineal region.¹ They are malignant tumours that infiltrate the adjacent structures by direct invasion and tend to spread through cerebrospinal fluid (CSF) circulation.² MRI helps in depicting the local infiltration and leptomeningeal spread of the disease. Tumour spread occurs as a result of CSF seeding from direct extension of the tumour through the pia or ependymal lining.

We present an MRI of a 2-year-old boy, a biopsy-proven case of pineoblastoma, which revealed a large lobulated mass in the pineal region. It was iso to hypointense on T1-weighted images (T1WIs) (figure 1A) and heterogeneously hyperintense on T2-weighted images (figure 1B). It showed restricted diffusion, seen as hyperintensity on

diffusion-weighted image (figure 1C) and hypointensity on corresponding apparent diffusion coefficient image (figure 1D). It showed heterogeneous enhancement on post-gadolinium T1WIs (figure 1E). It was infiltrating the tectum, cerebral aqueduct and third ventricle, and also extending into the fourth ventricle. Secondary obstructive hydrocephalus was present (figure 2A). Leptomeningeal spread was seen as extensive areas of nodular enhancement on contrast images along the sulcal spaces in the cerebral hemispheres, bilateral sylvian fissures, suprasellar cisterns, cerebellar folia, perimesencephalic cisterns and ventricular wall (figure 2B and C). There were also extensive leptomeningeal metastases along the entire spinal cord, which were seen as nodular enhancement along the spinal cord (figure 2D).

The patient received radiotherapy and is doing well at 3 months of follow-up.

Figure 1 MRI showing a large lobulated mass in the pineal region. (A) Axial T1-weighted image (T1WI) MRI shows isointense to hypointense mass. (B) Axial T2-weighted image shows heterogeneously hyperintensity in the mass. Diffusion-weighted image and apparent diffusion coefficient images reveal restricted diffusion (C and D). (E) Post-gadolinium axial T1WI showing heterogeneous enhancement.

To cite: Sandip S, Das CJ, Khandelwal RK. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-210343

Figure 2 (A) Axial fluid-attenuated inversion recovery image showing hydrocephalus and nodularity along the wall of ventricles (white arrow). Post-gadolinium axial and sagittal T1-weighted images (T1WIs) reveal extensive areas of nodular enhancement along the sulcal spaces and cisterns (white arrows) (B and C). (D) Post-gadolinium sagittal T1WI of cervical spine shows nodular enhancement along the spinal cord.

Learning points

- ▶ Although diagnosis of leptomeningeal metastases of tumour is performed by positive cerebrospinal fluid (CSF) analysis, positive cytological findings are seen in only 15–60% cases of leptomeningeal metastases.³
- ▶ Hence, a negative CSF analysis does not exclude the possibility of leptomeningeal spread of disease against a background of strong clinical suspicion.
- ▶ Gadolinium diethyle netriamine penta acetic contrast-enhanced MRI has a high detection rate for leptomeningeal metastases compared to non-contrast MRI.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Gasparetto EL, Cruz LCH Jr, Doring TM, *et al.* Diffusion-weighted MR images and pineoblastoma: diagnosis and follow-up. *Arg Neuropsiquiatr* 2008; 66:64–8.
- 2 Reis F, Faria AV, Zanardi VA, *et al.* Neuroimaging in pineal tumors. *J Neuroimaging* 2006;16:52–8.
- 3 North C, Segall HD, Stanley P, *et al.* Early CT detection of intracranial seeding from medulloblastoma. *AJNR Am J Neuroradiol* 1985;6:11–13.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow