Penile epidermal inclusion cyst: a rare location

Parthasarathy Athmarao Ealai,¹ Vikas Kapildeo Yadav,¹ Harshad Arvind Vanjare,² Sridhar Gibikote²

¹Department of Radiology, Christian Medical College, Vellore, Tamil Nadu, India
²Christian Medical College, Vellore, Tamil Nadu, India

Correspondence to
 Dr Vikas Kapildeo Yadav,
 vkyadav77@yahoo.co.in

Accepted 5 August 2015

DESCRIPTION

A 12-year-old boy presented with painless, slowly progressive swelling over the dorsum of the penis, evolving over 6 years. The swelling was present just underneath the penile skin. It was mildly compressible without any signs of inflammation. There were no other systemic symptoms. The only remarkable history was of circumcision 9 years prior. The patient was referred for ultrasound and Doppler evaluation of the swelling with the suspicion of a vascular malformation. Ultrasound scan revealed a well-defined lesion beneath the dermis, with fine internal echoes within it. No vascularity was noted on Doppler interrogation (figure 1). A provisional diagnosis of epidermal inclusion cyst was made based on surgical history and typical ultrasound appearance.

Intraoperatively, the lesion was found underneath the dartos fascia and adherent to urethra. Histopathology was suggestive of epidermal inclusion cyst (figure 2). It revealed a fibrocollagenous cyst lined by stratified squamous epithelium with keratin inside. Adjacent stroma was infiltrated by lymphocytes and foreign body giant cells.

Various theories for formation of cysts include sequestration of epidermal rests during embryonic life, occlusion of pilosebaceous units and traumatic or surgical implantation of epithelial elements. The condition is rare in the penis and can occur after an apparently minor procedure such as circumcision.¹ In our case, an ultrasound scan showed homogenous internal echoes with absence of vascularity. Posterior acoustic enhancement with attachment to dermis are other ultrasound features.² The only

Figure 2 Histopathology section of the cyst wall with lumen. The cyst wall is lined with stratified squamous epithelium (white arrow) with the inner wall showing a granular layer (arrowhead). Central lumen showing keratin accumulation (black arrow).

definitive treatment is complete surgical excision without rupturing the cyst, to avoid recurrence.³

Learning points

- ▶ Epidermal inclusion cyst must be considered in the differential diagnoses of painless swelling over the penis in the proper clinical setting, especially in the paediatric age group with a history of circumcision.
- ▶ Characteristic ultrasound appearance of cystic lesion with homogenous internal echoes, posterior acoustic enhancement with attachment to dermis and absence of vascularity should help diagnosing this entity.
- ▶ Delayed complication of epidermal inclusion cyst must be borne in mind in any postoperative site.

Acknowledgements The authors wish to thank Dr Mayank Gupta, Department of Pathology, Christian Medical College, for the histopathology image.

Contributors PAE contributed to drafting and designing the manuscript. VKY contributed to drafting, analysing and revising the manuscript critically for important intellectual content and for final approval. HAV contributed to acquisition and interpretation of the data. SG contributed to revising the manuscript critically for important intellectual content and for final approval.

Competing interests None declared.

Patient consent Obtained.

Figure 1 Well-defined lesion over the dorsum of the penis underneath the dermis, with fine internal echoes and posterior acoustic enhancement (white arrow on the left). No vascularity is seen within the lesion on Doppler (right side).

To cite: Ealai PA, Yadav VK, Vanjare HA, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-212137

Provenance and peer review Not commissioned; externally peer reviewed

REFERENCES

- 1 Saini P, Mansoor MN, Jalali S, *et al.* Penile epidermal inclusion cyst. *Indian J Pediatr* 2010;77:815–16.
- 2 Huang CC, Ko SF, Huang HY, *et al.* Epidermal cysts in the superficial soft tissue sonographic features with an emphasis on the pseudotestis pattern. *J Ultrasound Med* 2011;30:11–17.
- 3 Park HJ, Park NC, Park SW, *et al.* Penile epidermal inclusion cyst: a late complication of penile girth enhancement surgery. *J Sex Med* 2008;5:2238–40.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow