'Dilated odontoma': an incidental finding

Ali Zakir Syed,¹ Aravind Parachuru Venkata,² Rui Amaral Mendes³

¹Department of Oral and Maxillofacial Medicine and Diagnostic Sciences, Case School of Dental Medicine, Cleveland, Ohio, USA

²Department of Paediatric Dentistry, School of Dentistry, Dunedin, New Zealand

³Department of Oral and Maxillofacial Medicine and Diagnostic Sciences, Case Western Reserve University, Cleveland, Ohio, USA

Correspondence to

Dr Ali Zakir Syed,
azs16@case.edu

Accepted 25 September 2015

DESCRIPTION

The authors report an incidental finding in a 23-year-old man: an unusual case of dens invaginatus (type III), in relation to an upper left maxillary tooth (tooth #28). The patient presented for a routine dental check-up to the department of oral and maxillofacial medicine and diagnostic sciences. No significant medical history was elicited. Panoramic and full-mouth radiographic series were obtained after a detailed clinical examination (figures 1 and 2).

This rare malformation usually occurs in primary, permanent or supernumerary teeth.¹ Although frequently and erroneously called 'dilated odontoma', dens invaginatus must be differentiated from an odontoma; it is a well-known and described complex variant. Dens invaginatus should not be regarded as a tumour per se, but rather as a tooth malformation that often presents itself as a single calcified structure with a more radiolucent central portion.¹⁻³

The correct terminology for this malformation should be dens in dente or dens invaginatus type III. The term 'dilated odontoma' was first coined to describe a severe kind of dens invaginatus and must not be considered as a type of odontoma, which is a hamartomatous tumour, according to the WHO's Classification of Head and Neck Tumours.²

The absence of additional dental abnormalities in the current case can be confirmed in the panoramic dental radiograph (figure 2), which helps to preclude other abnormalities that might lead to consideration of potential genetic causes involving genes known to participate in the development of teeth, particularly the following five: MSX1, DLX1 and 2, PAX9 or PITX2.¹⁻³

Although type III dens invaginatus often presents as an invagination extending through the root, with enamel lining the invagination, the anatomical location, orientation and type of radiographic examination routinely performed during the first

Figure 2 Panoramic radiograph showing dilated odontoma with radiolucent centre and radiopaque periphery giving an appearance of doughnut.

appointment did not enable us to assess such a feature. Advanced imaging such as cone beam CT could have been performed. However, the cost involved and lack of impact on diagnosis and management precluded us from carrying on any further advanced imaging.⁴

The patient was referred to the oral and maxillofacial surgery department for surgical evaluation and management.

Learning points

- ▶ It is important to recognise 'dilated odontoma' as a dental anomaly rather than a true hamartoma.
- ▶ The anomaly has the characteristic radiographic feature of a radiolucent centre surrounded by radiopaque entity presenting as a doughnut shape.
- ▶ Combining ancillary imaging methods with a thorough clinical history is mandatory to attain an adequate diagnosis and for subsequent management.

Competing interests None declared.

Patient consent Not obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Sedano HO, Ocampo-Acosta F, Naranjo-Corona RI, et al. Multiple dens invaginatus, mulberry molar and conical teeth. Case report and genetic considerations. *Med Oral Patol Oral Cir Bucal* 2009;14: E69-72.
- 2 Takeda Y, Tomich CE. Ameloblastic fibro-odontoma. In: Barnes L, Eveson JW, Reichart P, et al, eds. *Pathology and genetics of head and neck tumours*. Lyon: IARC Press, 2005:283-327.
- 3 Chhabra N, Goswami M, Chhabra A. Genetic basis of dental agenesis—molecular genetics patterning clinical dentistry. *Med Oral Patol Oral Cir Bucal* 2014;19:e112-19.
- 4 Pradeep K, Charlie M, Kuttappa MA, et al. Conservative management of type III dens in dente using cone beam computed tomography. *J Clin Imaging Sci* 2012;2:5.

Figure 1 Intraoral periapical radiograph demonstrating dilated odontoma in the upper left maxillary projection.

CrossMark

To cite: Syed AZ, Parachuru Venkata A, Mendes RA. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-212594

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow