Spots of bother

Samantha Mary-Anne Morgan,¹ Barnaby Flower,² Mark A Samaan³

¹Barnet General Hospital, London, UK

²Newham University Hospital, London, UK

³Department of Gastroenterology, University College London Hospital, London, UK

Correspondence to

Dr Mark A Samaan, markasamaan@gmail.com

Accepted 28 November 2015

DESCRIPTION

A 24-year-old man presented with a 2-day history of a purpuric rash affecting his lower limbs, associated with fever, malaise and anorexia. He had fistulising, stricturing Crohn's disease with an end ileostomy and two enterocutaneous fistulae requiring long-term parenteral nutrition (PN) via a single-lumen Hickman line. His only medication was analgesia and vitamin B₁₂. He had not received regular steroids or immunosuppressive therapy for over 2 years.

On examination, he was febrile and tachycardic but not hypotensive. There were no cardiac murmurs and chest auscultation was unremarkable. His abdomen was soft and stoma output unchanged. A purpuric rash was visible on his lower limbs (figure 1). There was no meningism and examination of eyes, ears and throat was unremarkable.

Blood tests revealed haemoglobin 15.2 g/dL, white cell count 11.5 (neutrophils 9.5, eosinophils 0.8) $\times 10^6$ /mL, platelet count 247×10^9 /L, erythrocyte sedimentation rate 14 mm/h, C reactive protein 82.1 mg/L, sodium 122 mmol/L, potassium 4.6 mmol/L, urea 13.8 mmol/L, creatinine 89 μ mol/L, bilirubin 21 μ mol/L, albumin 28 g/L, alanine transaminase 133 U/L, alkaline phosphatase 673 U/L, international normalised ratio 1.4 and activated partial thromboplastin time 42 s. Chest X-ray and urinalysis were normal. An ultrasound and CT scan of the abdomen showed a

fatty liver consistent with PN-associated liver disease but no other abnormalities.

Candida glabrata was isolated from Hickman line blood cultures after 24 h incubation. The line was removed and 2 weeks of intravenous caspofungin started. The rash resolved within 48 h of treatment and inflammatory markers normalised. A transthoracic echocardiogram, vasculitis screen and complement levels were normal.

Learning points

- ▶ Parenteral nutrition is an important risk factor for invasive candidiasis; *Candida glabrata* accounts for approximately 17% of positive cultures. Clinicians should remain alert to the possibility of disseminated candidiasis in susceptible patients.
- ▶ The appearance of skin lesions can be the first clinical manifestation of acute disseminated candidiasis and this diagnosis should be considered in susceptible patients presenting in this manner.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

CrossMark

To cite: Morgan SM-A, Flower B, Samaan MA. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-212721

Purpuric rash

48 hours after treatment with Caspofungin

Figure 1 Purpuric rash pretreatment and post-treatment.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow