Correcting hypotension: be careful!

Ellis Foran, Tarig Mohammed Abkur

Department of Neurology, HSE
West, Galway, Ireland

Correspondence to

Dr Tarig Mohammed Abkur,
tarig1982@hotmail.com

Accepted 25 September 2015

DESCRIPTION

A 42-year-old woman presented with a 2-day history of vomiting. She was drowsy, dehydrated, tachycardic and hypotensive with a blood pressure (BP) of 52/37. The immediate management included infusion of 4 L of normal saline and inotropic support. Investigations showed elevated creatinine at 190 mmol/L, neutrophil leucocytosis and hypomagnesaemia at 0.55 mmol/L.

The patient was started on piperacillin/tazobactam for treatment of urinary tract infection. Her BP increased to 155/105 after 8 h.

On day 10 of admission, our patient reported headache, became confused and developed generalised tonic clonic seizure. Urgent CT of the brain showed bilateral posterior hypodensity (figure 1).

A diagnosis of posterior reversible encephalopathy syndrome (PRES) was suspected based on clinical grounds and was further supported by the radiological appearance on CT.

Findings on MRI of the brain were typical of PRES, with bilateral hyperintensities involving posterior parietal and occipital lobes (figure 2). These changes were in a similar distribution to the abnormalities seen on the CT. There was no evidence of restricted diffusion on diffusion-weighted imaging.

The patient improved and repeat MRI 3 months later showed resolution of the previously described changes.

PRES is a reversible neurological syndrome initially described by Hinchey *et al.*¹ It is characterised by the presence of white matter oedema affecting mainly the parieto-occipital region, although involvement of other areas of the brain has been reported. Hypertensive encephalopathy is the most

Figure 1 Axial non-contrast CT of the brain showing bilateral posterior hypodensity (see arrows).

Learning points

- ▶ PRES is a clinoradiological syndrome characterised by headache, seizures and visual changes, along with typical imaging features, primarily in the posterior regions of the brain.
- ▶ The rate of increase in blood pressure (BP) is a more important factor in the development of PRES than the absolute BP level.
- ▶ Hypomagnesaemia is considered a risk factor for PRES due to fact that it increases the permeability of the blood–brain barrier.

Figure 2 (A) Axial fluid-attenuated inversion recovery (FLAIR) MRI. (B) Sagittal FLAIR MRI showing hyperintensities affecting the posterior parietal and occipital lobes (see arrows). (C) Diffusion-weighted image showing no evidence of infarction.

CrossMark

To cite: Foran E, Abkur TM.
BMJ Case Rep Published
online: [please include Day
Month Year] doi:10.1136/
bcr-2015-212904

common cause and the rate of BP rise is more important than the absolute BP level.^{2 3}

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Hinchey J, Chaves C, Appignani B, *et al.* A reversible posterior leukoencephalopathy syndrome. *N Engl J Med* 1996;334:494–500.
- 2 Marik PE, Varon J. Hypertensive crises: challenges and management. *Chest* 2007;131:1949–62.
- 3 Vaughan CJ, Delanty N. Hypertensive emergencies. *Lancet* 2000;356:411–17.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow